

Review

A Brief Review on Engineering Synthetic Regulators for Bi-Directional Gene Expression Regulation

Thales Spartalis and Xun Tang *

Cain Department of Chemical Engineering, Louisiana State University, Baton Rouge, LA 70803, USA; tspart1@lsu.edu (T.S.)

* Corresponding author. E-mail: xuntang@lsu.edu (X.T.)

Received: 19 August 2026; Revised: 15 September 2026; Accepted: 15 October 2026; Available online: 27 October 2026

ABSTRACT: Since the advent of the first synthetic gene circuit, the field of gene circuits design has made great progress over the past several decades, bringing in feedback controllers, biomolecular neural networks, *etc.* The foundation of these circuits lies in their capability to precisely regulate gene expression in response to environmental cues. In this brief review, we focus on the constituent gene activation and repression regulatory parts, discuss their application and current progresses in bi-directional gene expression regulations, given the critical importance of such capability in ensuring a wide range of circuit properties, such as reference tracking, disturbance rejection, and system stability. We also discuss the integration of mathematical modeling in facilitating circuits design to complement the review. We anticipate more efforts to develop synthetic gene regulators that can repeatedly up- and down-regulate target gene expression dynamics in the future.

Keywords: Gene circuits; Bi-directional gene regulation; CRISPR; Riboregulator; Control; Modeling

1. Introduction

Precise control of protein expression is critical in maintaining healthy function in biosystems, as protein misfolding and abnormal protein homeostasis directly contribute to the pathogenesis of numerous diseases, including cancer, metabolic disorders, and neurodegenerative conditions [1–3]. The ability to achieve such precise modulation of gene expression is thus a cornerstone of synthetic biology, necessary for the development of novel diagnostics, therapeutics, and sustainable biomanufacturing processes [4–6]. One underdeveloped capability underpinning this objective is to achieve dynamic, reversible, and predictable control over gene expression, with both up- and down-regulation of the target gene, in response to specific signals or environmental cues [7]. This capability is essential for creating robust and adaptable biological systems that can function effectively in complex and changing environments. To meet this challenge, synthetic biologists have focused on developing gene circuits for engineering sophisticated dynamics. Note that while efforts have been made on both single and multiple gene expression regulation, in this review, we focus primarily on single gene expression regulation.

Synthetic gene circuits are composed of genetic parts that can regulate gene expression at transcriptional, translational, and post-translational levels [8] and can be designed to perform complex

computations, implement feedback control, and execute dynamic genetic programs [9,10]. The pioneer synthetic gene circuits, the genetic toggle switch, are constructed from two repressible arcs arranged in a mutually inhibitory network, demonstrating the feasibility of engineering bistable systems, but lacked the ability for graded and reversible control [11,12]. Over the past several decades, the complexity of gene circuits has grown exponentially, bringing in multi-layered architectures, logic gates, and intricate feedback loops [7,13].

One class of gene circuits that is well-known for maintaining target gene expression in the presence of disturbance is the antithetic controller, a negative feedback-based gene circuit that leverages molecular sequestration to realize bi-directional regulation. The concept of feedback is central to biological systems, where positive feedback loops can create switch-like behaviors and amplify cellular responses [14,15], and negative feedback loops are essential for maintaining stable protein concentrations and preventing overshoots [16,17]. Inspired by these natural control principles, synthetic biologists have implemented sophisticated feedback strategies in engineered microbial consortia. For instance, Aoki et al. (2019) developed a universal biomolecular integral feedback controller in *E. coli*, inspired by an antithetic feedback motif, that mathematically ensures robust perfect adaptation, meaning that the system can maintain the target gene expression at the setpoint despite disturbances [18]. Subsequent efforts extended this concept to multicellular systems, in which control functions are distributed across different cell populations to reduce metabolic burden and enhance modularity. In Martinelli et al. (2022), the authors designed a multicellular Proportional-Integral controller where proportional and integral actions were distributed across two different cell populations, and communicated via orthogonal quorum sensing molecules to regulate a third target population [19]. By distributing the sense, actuate, and control functions across different cell populations, these multicellular systems can achieve remarkable precision and stability in gene expression, thereby creating closed-loop feedback systems that enhance circuit robustness and enable the regulation of complex, multi-cellular behaviors.

Instead of focusing on specific types of controllers, such as the antithetic controller described above, this review focuses on key technologies that enable up- and down-regulation of single-gene expression for bi-directional regulation, such as CRISPR-based systems and RNA-based regulatory mechanisms. To complete the discussion, we also examine the application of mathematical modeling to guide the design and characterization of these circuits.

2. RNA-Based Gene Expression Repression and Activation Regulators

Living organisms have evolved intricate regulatory networks that maintain cellular homeostasis and enable adaptation to changing environments through precise control of gene expression in response to internal and external cues [18,20–25]. In bacteria, post-transcriptional regulation is achieved through a diverse class of molecules known as regulatory RNAs [26–28]. These molecules are commonly grouped into classes according to their mechanism and layer of action: antisense and small RNAs, which base-pair with a target mRNA to repress translation or promote its degradation; small transcription activating RNAs (STARs), which relieve transcription termination to switch on a downstream gene; and toehold switches and related riboregulators, which expose a sequestered ribosome binding site upon binding a trigger RNA to activate translation [26]. These regulatory RNAs traditionally function by targeting mRNAs to induce degradation or suppress translation [27,29,30]. This natural mechanism has provided a foundational blueprint for synthetic biology, where engineered regulatory RNAs with predictable behavior are now routinely used to optimize metabolic pathways, down-regulate competing gene expression, and perform targeted gene knockdowns in complex systems [31]. As a class, these antisense and small regulatory RNAs constitute the repression arm of the RNA toolbox, acting post-transcriptionally and imposing minimum metabolic load because they are encoded as short, non-translated sequences [32,33].

Efforts over the past several decades have expanded the regulatory paradigm beyond gene repression to synthetic modes of activation. Small transcription activating RNAs (STARs), for instance, can be engineered to disrupt intrinsic transcription terminators, leading to robust gene activation, enabling the construction of RNA-only transcriptional logic gates. This evolution from repurposing natural components to forward-engineering novel functions provides a rich, dynamic toolbox for achieving programmable bi-directional control [26,34–36]. Besides transcriptional regulation, RNA-based regulators can also operate at the post-transcriptional level, controlling translation initiation or termination. One prominent example is the “toehold switch”, a de-novo-designed riboregulator that can activate gene expression in the presence of a specific trigger RNA [37,38]. These switches exhibit a wide range of dynamics with orthogonality, meaning that they can be designed to respond to unique RNA sequences with minimal crosstalk [39,40]. This capability permits independent regulation of multiple genes within the same cell, as demonstrated by the construction of a four-input AND logic gate [13]. The design principles underlying toehold switches have been further refined, with recent work focusing on optimizing the thermodynamic properties of the RNA structures to maximize dynamic range and minimize leakiness [4,40–42]. McSweeney et al. [36] explored the use of short activators and repressors of RNA toehold switches, demonstrating that the length and sequence of the trigger RNA can significantly impact the performance of the switch. Considered together, these tools span complementary layers of control, with antisense and small RNAs repressing post-transcriptional activities, STARs activating during transcription elongation, and toehold switches activating at translation initiation, so that bidirectional regulation at the RNA level is generally realized by pairing an activating and a repressing device rather than by a single part [27,35,37].

As compared to protein-based regulators, these RNA-based devices can offer a complementary and highly programmable approach to gene regulation [26,43]. One example is the combined use of STARs and antisense RNAs [28,31,34,44]. The STAR system provides a method for activating gene expression, which can be combined with traditional antisense RNA-mediated repression to achieve multilevel control, as shown by Lee and Moon [35]. These approaches provide a fast and versatile means of regulation that does not require the expression of bulky protein factors, making them ideal for applications where a rapid response is needed [2]. Their principal limitations, relative to protein- and CRISPR-based regulators, are a susceptibility to incomplete repression or leakage and a reliance on predictable RNA folding, which can restrict dynamic range and portability across host contexts [31,42]. These trade-offs, and how they compare with the protein- and CRISPR-based platforms introduced in the following sections, are discussed in greater detail later in the manuscript.

The scope of RNA-based regulation has been further extended by Simmons et al. [45], who rewired native post-transcriptional global regulators to construct designer, multi-layered genetic circuits, demonstrating that endogenous RNA regulatory networks can be repurposed as programmable circuit components. The use of antisense RNAs for gene regulation has a long history in synthetic biology, and recent work has focused on improving the predictability and efficiency of these systems [29,31,33,38,46,47]. By carefully designing the complementary parts between the antisense RNA and its target mRNA, it is possible to achieve precise control over translation and mRNA stability [31,39].

Another powerful class of RNA-based or RNA-protein hybrid regulators is the CRISPR-Cas system, originally discovered as a bacterial immune system, which has been repurposed into a versatile tool for programmable gene regulation [48]. The most widely used effector, Cas9 from *Streptococcus pyogenes*, is an RNA-guided endonuclease directed by a single-guide RNA (sgRNA) to a complementary genomic sequence adjacent to a protospacer-adjacent motif (PAM), where it introduces a double-strand break. Catalytic inactivation of its two nuclease domains produces a nuclease-dead variant, dCas9, which retains guide-directed DNA binding but no longer cleaves the target sequence [48,49]. Fusion of dCas9 to transcriptional repressor domains, such as KRAB, or activator domains, such as VP64 or the tripartite VPR complex, converts sequence-specific DNA targeting into programmable gene repression (CRISPR

interference, CRISPRi) or activation (CRISPR activation, CRISPRa), respectively [49,50]. The versatility of CRISPR-based tools for controlling gene expression in bacteria has been comprehensively reviewed by Vigouroux and Bikard [6], who highlighted the diversity of Cas proteins and the different mechanisms by which they can be harnessed for synthetic biology applications, from transcriptional regulation to post-transcriptional control. One breakthrough that leverages this versatility is the development of multilayer CRISPRa/i circuits. For instance, Tickman et al. [7] demonstrated that these circuits can operate effectively in both cell-free and bacterial systems, enabling dynamic genetic programs with unprecedented complexity. Their work showed that by carefully tuning the strength of guide RNAs and the architecture of activatable promoters, it is possible to create circuits with multiple layers of regulation, in which the output of one layer serves as the input to the next. This hierarchical organization is reminiscent of natural gene regulatory networks and provides a powerful framework for engineering complex behaviors. In Nielsen and Voigt [51], the authors also provided a foundational framework by building a set of orthogonal dCas9-based NOT gates in *E. coli*. These repressive modules were layered to create more complex logic, and by targeting the native transcription factor *maltT*, they showed that synthetic circuits could reprogram complex cellular phenotypes, establishing a principle for scalable, programmable genetic control.

Besides independent usage, these RNA-based devices can also be coupled with other regulatory mechanisms for new functions. Hong et al. [4] demonstrated the use of multilevel gene regulation using switchable transcription terminators and toehold switches in *E. coli*. This approach enabled independent control of transcription and translation, providing a powerful framework for engineering complex regulatory networks. Similarly, Zhao et al. [52] demonstrated the construction of a three-layer cascade circuit and a two-input, three-layer OR logic gate using switchable transcription terminators in a cell-free system, highlighting the potential to create complex, multi-layered circuits. The use of RNA scaffolds has also emerged as a promising strategy for organizing metabolic enzymes and enhancing pathway flux. Mitkas et al. [53] and Pham et al. [46] showed that Cas6-mediated RNA scaffolding can be used to assemble metabolons in yeast, demonstrating the potential of RNA-based approaches for metabolic engineering. Furthermore, Xi et al. [54] demonstrated that guide RNAs can be conditionally deactivated by mRNA and small molecule triggers in *S. cerevisiae*, providing a strategy for integrating RNA-sensing with CRISPR-based gene regulation. The integration of CRISPR-based regulation with riboswitches and STARs provides a rich palette of tools for engineering sophisticated genetic circuits [34,55]. Figure 1 provides an overview of the aforementioned RNA-based regulators, possible design strategies and applications.




Figure 1. Molecular strategies for bi-directional gene expression control and their applications in synthetic biology. The figure illustrates the two primary molecular platforms for achieving simultaneous up- and down-regulation of gene expression:

CRISPR-based systems, in which a catalytically deactivated Cas9 (dCas9) protein is directed by guide RNAs to activate (CRISPRa) or repress (CRISPRi) transcription at specific genomic loci, and RNA-based mechanisms, including toehold switches, antisense RNAs, and STARS, which operate at the post-transcriptional level. Both platforms can be assembled into multilayer logic circuits (AND, OR, NOT gates) for complex regulatory programs. Natural regulatory networks, which achieve homeostasis and adaptation through feedback control, serve as the biological inspiration for these engineered architectures. Mathematical modeling approaches, including ODE-based models and control theory frameworks, provide the computational tools necessary for rational design and performance prediction of these circuits. Together, these strategies find application across diagnostics, therapeutics, environmental sensing, metabolic engineering, and biomanufacturing.

3. Protein-Based Gene Expression Repression and Activation Regulators

Beyond the RNA-based devices discussed in the previous section, which offer high design flexibility and comparatively low metabolic burden [32], protein-based transcription factors have played a central role in synthetic gene regulation since the inception of the field, as exemplified by foundational circuits such as the genetic toggle switch and the repressilator [12,56,57]. However, the repertoire of well-characterized natural regulators is limited, and the simultaneous use of multiple regulators can introduce undesirable crosstalk [58]. To address these limitations, programmable transcriptional regulators have been engineered from modular DNA-binding proteins. Prominent examples include zinc-finger and transcription activator-like effector (TALE) domains fused to activation or repression modules, yielding orthogonal and tunable regulators for both bacterial and eukaryotic systems [59–61]. More recently, this strategy has been extended toward therapeutic applications through compact, largely human-derived zinc-finger regulators that enable small molecule-controlled expression of therapeutic genes in primary human T cells [62].

A distinct strategy in bacteria exploits cognate sigma/anti-sigma factor pairs. In this architecture, an alternative sigma factor recruits RNA polymerase to a cognate promoter to activate transcription, whereas the corresponding anti-sigma factor sequesters the sigma factor to repress transcription. This sequestration-based mechanism is conceptually reminiscent of the antithetic controllers discussed previously. Rhodius et al. (2013) systematically characterized 86 extracytoplasmic function (ECF) sigma factors and 62 anti-sigma factors, identifying mutually orthogonal regulator pairs that can function as programmable genetic switches in *E. coli* [63].

Regulation can also be directed at the transcriptional machinery itself. Bacteriophage T7 RNA polymerase, when split into interdependent fragments, can reconstitute an orthogonal and largely host-independent transcription system in which a shared “core” fragment modulates the overall transcriptional capacity of the circuit. This architecture has been exploited to construct tunable “resource allocators” and orthogonal AND-gate inputs [64,65], and subsequent directed-evolution efforts have expanded the utility of these orthogonal transcription systems for programmable gene expression in both prokaryotic and eukaryotic hosts [66]. Whereas transcriptional and translational regulation primarily control the rate of protein synthesis, the decline of an existing protein pool is governed by dilution and degradation. Active, targeted proteolysis has therefore been used to accelerate protein down-regulation [67,68]. Building on engineered *ssrA* degradation tags and redesigned protease-recognition sequences [69,70], inducible and orthogonal degradation systems enable the steady-state abundance and active removal rate of target proteins to be tuned with a degree of independence [67,68]. More recent designs have extended inducible degradation to blue-light control in *E. coli* and to repurposed E3 ubiquitin ligase-based degron systems in mammalian cells [71,72]. By actively removing protein rather than merely halting its synthesis, controlled proteolysis is particularly well suited for rapid and reversible turn-off, a key requirement for symmetric bidirectional regulation.

These protein-based regulators differ fundamentally in the molecular layer at which they act and in the trade-offs they impose, as summarized together with other regulator classes in Table 1. Engineered transcription factors can provide high targeting specificity and support both activation and repression through selection of the effector domain, but they require expression of relatively large protein regulators

and may therefore increase cellular burden [58–61]. Sigma/anti-sigma systems are comparatively compact and readily compatible with feedback architectures, although their action is confined to transcription initiation [63]. Split polymerase systems can partially decouple circuit transcription from host machinery and make resource allocation an explicit design variable, directly linking them to competition for shared transcriptional resources discussed below [64–66]. Controlled proteolysis adds a post-translational regulatory layer that can accelerate down-regulation beyond what is typically achievable through repression of synthesis alone [67,68,71,72]. In this respect, controlled proteolysis and sequestration-based regulators are especially relevant to single-target control modes in which reversible turn-off over time is a primary requirement. In practice, these mechanisms are increasingly combined with one another and with RNA- and CRISPR-based regulators to support dynamic and symmetric up- and down-regulation for applications such as reference tracking, disturbance rejection, and stable bidirectional control [8,58].

4. Bi-Directional Gene Expression Regulations

The concept of combining activation and repression within a single genetic circuit has its roots in the study of natural network motifs, such as the incoherent feedforward loop (IFFL), where a single input simultaneously activates a target gene and its repressor. Bleris et al. [73] constructed synthetic IFFL circuits in human HEK293 cells, using a transcription factor to activate the expression of both a reporter gene and its post-transcription repressor microRNA. They found that this architecture can lead to adaptation to gene dosage, with the post-transcriptional miRNA-based version showing superior performance and lower noise. This work established the IFFL as a powerful motif for engineering robustness in synthetic circuits. Building on this, Guo and Murray [74] demonstrated the construction of modular IFFL circuits in an *E. coli* cell-free transcription-translation (TXTL) system and in living cells. Their circuit used a sigma factor to activate both a repressor (*TetR*) and a target gene (GFP). This study highlighted the power of cell-free systems for rapid prototyping, as the pulse-like dynamics observed *in vitro* closely matched the *in vivo* performance, validating the TXTL system as a reliable proxy for cellular behavior. These early synthetic implementations of the IFFL, while not constituting bidirectional regulation in the strict sense, established a foundational principle: that combining activation and repression within a single genetic circuit can produce functional, predictable outputs. The development of the aforementioned CRISPR-Cas technology and RNA-based devices provided the programmability and scalability that can potentially extend this principle into true multi-target, simultaneous up- and down-regulation of gene expression.

One prominent success is the combined use of the CRISPRi and CRISPRa systems. In Martella et al. [75], the authors systematically evaluated CRISPRa and CRISPRi modalities in human cells, identified dCas9-VPR (activation) and dCas9-KRAB (repression) as the most effective orthogonal pair, and demonstrated simultaneous up-regulation of one endogenous gene and down-regulation of another, showcasing the feasibility of bi-directional CRISPR regulation for controlling complex cellular processes in human cells. However, integrated use of CRISPRi/a systems in one host system is nontrivial, as crosstalk between components and off-target effects can complicate predictable circuit behavior [76]. As demonstrated by Barbier et al. [10], combining CRISPRi with a SoxS-based CRISPRa system in *E. coli* can lead to unpredictable, context-dependent effects due to different binding affinities of the guide RNAs for the activation and interference complexes. Their work highlighted the importance of component design, showing that a guide RNA with the same scaffold structure for both complexes can mitigate such issues and restore predictable circuit behavior. To further advance the scalability of these systems, Burbano et al. [9] engineered a suite of high-performing, activatable promoters with up to 1000-fold dynamic range, enabling CRISPRa/i circuits that are six layers deep and four branches wide, and demonstrating applicability for complex network topologies including feedback loops, logic gates, and dynamic pulse modulators. Most recently, Pacalin et al. [77] introduced CRISPRai, a system for simultaneous bidirectional

epigenetic editing in human primary T cells. Using two orthogonal dCas9 proteins fused to activation (*VPR*) and repression (*KRAB*) domains, they performed pooled screens coupled with single-cell RNA sequencing.

This allowed them to map epistatic relationships between genes by observing the transcriptional consequences of activating one gene while repressing another, revealing hierarchies in gene regulation and demonstrating the power of bi-directional perturbation as a tool for fundamental biological discovery. Recent advances have also focused on making CRISPR circuits more responsive to external signals. Kim et al. [78] developed modular and signal-responsive transcriptional regulation systems using CRISPRi-aided genetic switches in *E. coli*. These switches can be activated by specific small molecules or environmental cues, allowing for the dynamic control of gene expression in response to changing conditions. This capability is essential for applications in biosensing and adaptive biomanufacturing, where the ability to respond to environmental signals is critical.

The practical value of multidirectional regulation is particularly evident in metabolic engineering, where the design objective is often to redirect metabolic flux by up-regulating rate-limiting or committed pathway enzymes while simultaneously repressing competing or by-product-forming branches. This inherently bidirectional control problem can be addressed through complementary classes of regulators. Synthetic small regulatory RNAs, for example, have been used to tune pathway flux by post-transcriptionally suppressing competing genes [33], whereas CRISPR-based platforms enable simultaneous activation and repression of multiple pathway targets within a unified regulatory framework [79]. Lian et al. [79] introduced CRISPR-AID, a tri-functional CRISPR system in *Saccharomyces cerevisiae* that combined CRISPRa, CRISPRi, and gene knockout in a single framework. Applied to the β -carotene biosynthesis pathway, they achieved a 3-fold increase in production by simultaneously activating three genes, repressing one, and deleting another, showcasing the power of this approach for combinatorial metabolic engineering. Chen et al. [80] applied simultaneous CRISPRa and CRISPRi to improve the performance of the electroactive bacterium *Shewanella oneidensis*. They activated genes to increase cell length and biofilm formation while repressing a negative regulator of flavin synthesis. This combined perturbation led to a 2.5-fold increase in power density in a microbial fuel cell, illustrating how coordinated multi-directional regulation can optimize complex, multi-trait phenotypes. Extending this to the industrial yeast *Yarrowia lipolytica*, Chen et al. [81] further developed a CRISPR-STAR platfo that uses a single dCas9 directed by scaffold RNAs (scRNAs) to recruit either an activator (*VPR*) or a repressor (*MXI1*). This simplified architecture was used to enhance fatty alcohol production by simultaneously activating a key enzyme and repressing a competing pathway, leading to a 55.7% increase in yield and demonstrating the effectiveness of this platform for metabolic optimization.

Taken together, the systems discussed in this section illustrate how combined activation and repression have evolved from early single-target motifs into programmable platforms capable of regulating multiple genes simultaneously. CRISPR-based tools, in particular, now enable coordinated up- and down-regulation of multiple endogenous targets for applications ranging from epistatic mapping to metabolic engineering. These applications exemplify the multi-target form of bidirectional regulation introduced earlier, which is distinct from reversible control of a single target over time, although the two modes can be combined in practice. As summarized in Table 1, the regulator classes reviewed here generally offer complementary strengths rather than a single universally superior solution: RNA-based devices are highly programmable and can impose relatively low burden, although individual designs may exhibit leakage [31,32]; protein-based regulators can provide broad dynamic range and intrinsic bidirectionality at the cost of greater cellular burden [58,59]; and CRISPR-based systems offer programmable multiplexing while remaining dependent on shared cellular resources [49,82].

Table 1. Comparison of synthetic regulator classes for bidirectional gene expression control.

Regulator Class	Regulatory Layer (Mechanism)	Direction	Dynamic Range	Orthogonality and Multiplexing	Metabolic Burden	Programmability	Representative Examples
Antisense and small regulatory RNAs	Post-transcriptional; base-pairs with mRNA to block translation or trigger degradation	Repression	Set by base-pairing; improved by design rules, not standardized across studies [31]	High; determined by base-pairing sequence [31,33]	Low (short non-coding RNA, no protein product) [32]	High; retargeted by complementary sequence [31,33]	Pathway flux control, gene knockdown, antibacterial sensitization [31,33,47]
STARs (small transcription activating RNAs)	Transcriptional; disrupt intrinsic terminators (anti-termination)	Activation	Large; computationally optimized for high activation [34]	High; orthogonal libraries, multilevel with antisense RNA [34,35]	Low [32]	High; <i>de novo</i> sequence design [28,34]	RNA-only transcriptional logic, multilevel STAR and antisense control [28,34,35]
Toehold switches and riboregulators	Translational; trigger RNA exposes the ribosome binding site	Activation	Wide and tunable; leakiness reduced by thermodynamic optimization [4,40,42]	High, minimal crosstalk; four-input AND gate demonstrated [13,39]	Low [32]	Very high; <i>de novo</i> design [37,38]	Ribocomputing logic, native RNA detection, cell-free circuits [13,40,52]
CRISPRi and CRISPRa (dCas)	Transcriptional; dCas fused to repressor or activator effector	Both (CRISPRi represses, CRISPRa activates); orthogonal dCas9-KRAB and dCas9-VPR pair [75]	High; activatable promoters up to 1000-fold [9]	High targeting; six-layer, four-branch circuits [9]; constrained by the shared dCas pool and resource competition [82]	High (large dCas protein, draws shared resources) [82]	Very high; RNA-guided retargeting [48–50]	Bidirectional epigenetic editing [77]; metabolic engineering: 3-fold β -carotene [79], 2.5-fold power density [80], 55.7% fatty alcohol [81]; signal-responsive switches [78]
Protein transcription factors (classical and engineered ZF/TALE)	Transcriptional; sequence-specific DNA binding plus effector	Both (through effector choice)	High; broadly tunable [61]	Limited and crosstalk-prone for natural [58]; expanded by engineered ZF/TALE [59–61]	Moderate to high (expression of large protein factors) [58]	Moderate; modular ZF/TALE assembly but requires protein engineering [59,60]	Genetic toggle switch [12], synthetic IFFL [73], therapeutic T-cell circuits [62]
Sigma and anti-sigma factors	Transcriptional (initiation); redirect RNA polymerase	Both (sigma activates, anti-sigma represses by sequestration)	Moderate [63]	High; 86 ECF sigma and 62 anti-sigma factors mapped to orthogonal pairs [63]	Moderate (protein factors)	Low to moderate; requires cognate promoters [63]	Orthogonal genetic switches [63], IFFL activation module [74]

Split-T7 RNA polymerase	Transcriptional; orthogonal, largely host-independent polymerase	Activation and transcriptional-capacity control	High (strong T7 transcription)	Split mutants yield AND gates [64] and a tunable resource allocator [65] evolved orthogonal engines [66]	Moderate to high (possible T7 toxicity, resource draw) [65]	Low to moderate; T7 promoter variants [64]	Transcriptional AND logic [64], resource allocation [65], cross-host expression [66]
Controlled proteolysis and degrons	Post-translational; targeted protein degradation	Down-regulation (active protein removal)	Tunable degradation rate [67,68]	Few orthogonal protease and tag pairs; engineered recognition determinants [69,70]	Moderate (protease and degron expression)	Broad by tagging, but single direction [68]	Tunable degradation [67,68] light-inducible LOVdeg [71], E3-ligase degrons in mammalian cells [72]

5. Mathematical Modeling to Facilitate Circuits Design

The regulatory architectures described here, from RNA-based devices and CRISPR-mediated circuits to their applications in metabolic engineering, share a common challenge: as the number of interacting components increases, the behavior of the system becomes increasingly difficult to predict. Non-linear interactions between activators and repressors, competition for shared molecular resources such as *dCas9* or RNA polymerase, *etc.*, and the stochastic nature of gene expression at the single-cell level all contribute to emergent dynamics that cannot be reliably anticipated without a formal quantitative framework [10,82–84]. Mathematical modeling and computational simulation therefore serve not merely as complementary tools, but as necessary components of the design cycle, enabling researchers to explore the parameter space of a circuit, identify potential failure modes, and generate testable predictions before committing to the bench [85–87].

Predicting the dynamics of bi-directional gene circuits, which simultaneously employ activation and repression, presents a unique set of challenges, including capturing non-linear interactions, predicting emergent behaviors, and accounting for competition between opposing regulatory elements for shared molecular resources [10,82,88]. A purely empirical approach to circuit construction is often inefficient, costly, and susceptible to malfunction due to unexpected interactions between components, context-dependent behavior, and competition for limited cellular resources [82,86]. Mathematical modeling provides a rational framework for abstracting this complexity, enabling engineers to simulate circuit dynamics, perform sensitivity analyses to identify critical parameters, and optimize performance before experimental implementation [89]. Historically, the design of foundational synthetic circuits was tightly coupled with mathematical modeling. Ordinary differential equations (ODEs), which describe the rate of change of molecular concentrations over time, were used to define the parameter space in which landmark circuits like the genetic toggle switch and the repressilator were predicted to function [12,57]. This deterministic framework remains as the most popular type of modeling in synthetic biology due to its simplicity, providing a powerful tool for capturing the average behavior of a cell population. However, deterministic ODEs do not capture the stochastic fluctuations in gene expression that dominate at the single-cell level. Therefore, stochastic simulations emerged as alternative approaches to approximate these probabilistic dynamics [90–96]. Other studies subsequently demonstrated how to experimentally decouple cellular noise into two components: intrinsic noise arising from the inherent randomness of transcription and translation, and extrinsic noise stemming from fluctuations in the upstream cellular environment, such as variations in ribosome or polymerase concentrations [83,84,97]. The choice of modeling framework thus depends on the specific question, with deterministic models suited for population-level predictions and stochastic models essential for understanding noise propagation and cell-to-cell variability [98]. Beyond defining a circuit's operating regime, modeling can also specify the control architecture itself. Briat et al. [99] integrated stochastic chemical kinetics with control theory to show that a sequestration-based feedback motif can robustly maintain a target gene at its setpoint despite disturbances and intrinsic biochemical noise, thereby identifying a network topology capable of robust bidirectional regulation before experimental implementation. This theoretical architecture was subsequently realized as a synthetic integral feedback circuit in living cells, closing the loop between model-based design and experimental implementation [18].

As synthetic circuits grow in scale and complexity, particularly with the advent of multiplexed CRISPR and RNA-based regulation, models must account for the competition for shared cellular resources (e.g., *dCas9*, ribosomes, RNA polymerase) [86], and failure to do so leads to unintended crosstalk and breaks the modularity of circuit components. Model-guided design has been instrumental in addressing this challenge. For instance, a mathematical model of *dCas9* competition predicted that a negative feedback loop regulating *dCas9* expression could neutralize resource competition, a finding that was experimentally validated, restoring modularity in layered CRISPRi circuits [82]. More recently, Sechkar and Steel [88] used a similar

resource-aware model to show that a ‘push-pull’ architecture, where a gene is simultaneously activated and repressed, can confer robustness to resource competition crosstalk. The work further showed that combining positive and negative regulation in feedback loops can overcome trade-offs in robustness to different types of perturbations. Similarly, detailed ODE models have been developed for RNA-based circuits, such as toehold switches and STARs, enabling accurate prediction of their dynamic behavior and the model-guided discovery of previously unknown RNA maturation steps [89]. These models, often parameterized using data from experimental systems, are becoming increasingly sophisticated, forming the basis for computer-aided design tools that accelerate the engineering of complex, multi-layered regulatory networks [100,101]. Furthermore, besides the CRISPR and RNA-based systems, ODE models have also been used to characterize and optimize other bi-directional systems, such as light-activated and repressed promoters, revealing design principles for achieving fast and reversible switching dynamics [102]. These examples underscore the indispensable role of mathematical modeling in moving from the design of simple unidirectional switches to the engineering of sophisticated, robust, and predictable bi-directional regulatory systems [103].

6. Conclusions

The field of synthetic gene circuits has achieved significant breakthroughs over the past several decades, from the advent of the first repressilator to the latest biomolecular neural networks [104–107]. The underlying principle that enables the functionality of these circuits is the ability to precisely regulate the target gene dynamics. As the field progresses, increasing efforts are being made towards the construction of regulators that can repeatedly activate and repress target gene expression, to achieve properties analogous to mechanical feedback control systems, such as disturbance rejection, reference tracking, system stability, and robustness. The prosperity in the development of the constituent RNA-based regulatory parts such as STAR, riboregulators, and the CRISPR-Cas systems described in this review lays the cornerstone to the realization of advanced circuits for complicated functionalities with ensured properties. As the demand for precise and repeatable regulation of biological process increases, we expect bi-directional repetitive gene expression regulators to play an increasingly important role in future circuits design. Coupling with mathematical modeling and high throughput circuit characterization will be necessary for efficient and reliable design and construction.

Statement of the Use of Generative AI and AI-Assisted Technologies in the Writing Process

During the preparation of this manuscript, the authors used ChatGPT for language editing and references finding. After using this tool/service, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

Author Contributions

T.S.: conceptualization, investigation, writing—original draft, writing—review and editing. X.T.: conceptualization, funding acquisition, writing—original draft, writing—review and editing.

Ethics Statement

Not applicable.

Informed Consent Statement

Not applicable.

Data Availability Statement

No data were generated from this work.

Funding

This research was funded by NSF grant number #2223720.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Hetz C, Glimcher LH. Protein homeostasis networks in physiology and disease. *Curr. Opin. Cell Biol.* **2011**, *23*, 123–125. DOI:10.1016/j.ceb.2011.01.004
2. Gregersen N, Bross P, Vang S, Christensen JH. Protein Misfolding and Human Disease. *Annu. Rev. Genom. Hum. Genet.* **2006**, *7*, 103–124. DOI:10.1146/annurev.genom.7.080505.115737
3. Seo J, Park M. Molecular crosstalk between cancer and neurodegenerative diseases. *Cell. Mol. Life Sci.* **2020**, *77*, 2659–2680. DOI:10.1007/s00018-019-03428-3
4. Hong S, Kim J, Kim J. Multilevel Gene Regulation Using Switchable Transcription Terminator and Toehold Switch in *Escherichia coli*. *Appl. Sci.* **2021**, *11*, 4532. DOI:10.3390/app11104532
5. Morones-Ramírez JR. Biocircuitry and Living Programmable Materials: The Next Frontier in Synthetic Living Systems. *ACS Mater. Lett.* **2025**, *7*, 2910–2935. DOI:10.1021/acsmaterialslett.5c00238
6. Vigouroux A, Bikard D. CRISPR Tools To Control Gene Expression in Bacteria. *Microbiol. Mol. Biol. Rev.* **2020**, *84*, e00077-19. DOI:10.1128/MMBR.00077-19
7. Tickman BI, Burbano DA, Chavali VP, Kiattisewee C, Fontana J, Khakimzhan A, et al. Multi-layer CRISPRa/i circuits for dynamic genetic programs in cell-free and bacterial systems. *Cell Syst.* **2022**, *13*, 215–229.e8. DOI:10.1016/j.cels.2021.10.008
8. Buson F, Gao Y, Wang B. Genetic Parts and Enabling Tools for Biocircuit Design. *ACS Synth. Biol.* **2024**, *13*, 697–713. DOI:10.1021/acssynbio.3c00691
9. Alba Burbano D, Cardiff RAL, Tickman BI, Kiattisewee C, Maranas CJ, Zalatan JG, et al. Engineering activatable promoters for scalable and multi-input CRISPRa/i circuits. *Proc. Natl. Acad. Sci. USA* **2023**, *120*, e2220358120. DOI:10.1073/pnas.2220358120
10. Barbier I, Kusumawardhani H, Chauhan L, Harlapur PV, Jolly MK, Schaerli Y. Synthetic Gene Circuits Combining CRISPR Interference and CRISPR Activation in *E. coli*: Importance of Equal Guide RNA Binding Affinities to Avoid Context-Dependent Effects. *ACS Synth. Biol.* **2023**, *12*, 3064–3071. DOI:10.1021/acssynbio.3c00375
11. Fiore D, Salzano D, Cristobal-Coppulo E, Olm JM, di Bernardo M. Multicellular Feedback Control of a Genetic Toggle-Switch in Microbial Consortia. *IEEE Control Syst. Lett.* **2021**, *5*, 151–156. DOI:10.1109/LCSYS.2020.3000954
12. Gardner TS, Cantor CR, Collins JJ. Construction of a genetic toggle switch in *Escherichia coli*. *Nature* **2000**, *403*, 339–342. DOI:10.1038/35002131
13. Green AA, Kim J, Ma D, Silver PA, Collins JJ, Yin P. Complex cellular logic computation using ribocomputing devices. *Nature* **2017**, *548*, 117–121. DOI:10.1038/nature23271
14. Ferrell JE. Self-perpetuating states in signal transduction: Positive feedback, double-negative feedback and bistability. *Curr. Opin. Cell Biol.* **2002**, *14*, 140–148. DOI:10.1016/S0955-0674(02)00314-9
15. Cinquin O, Demongeot J. Roles of positive and negative feedback in biological systems. *Comptes Rendus. Biol.* **2002**, *325*, 1085–1095. DOI:10.1016/S1631-0691(02)01533-0
16. Yi TM, Huang Y, Simon MI, Doyle J. Robust perfect adaptation in bacterial chemotaxis through integral feedback control. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 4649–4653. DOI:10.1073/pnas.97.9.4649
17. Ma W, Trusina A, El-Samad H, Lim WA, Tang C. Defining Network Topologies that Can Achieve Biochemical Adaptation. *Cell* **2009**, *138*, 760–773. DOI:10.1016/j.cell.2009.06.013
18. Aoki SK, Lillacci G, Gupta A, Baumschlager A, Schweingruber D, Khammash M. A universal biomolecular integral feedback controller for robust perfect adaptation. *Nature* **2019**, *570*, 533–537. DOI:10.1038/s41586-019-1321-1

19. Martinelli V, Salzano D, Fiore D, di Bernardo M. Multicellular PI Control for Gene Regulation in Microbial Consortia. *IEEE Control Syst. Lett.* **2022**, *6*, 3373–3378. DOI:10.1109/LCSYS.2022.3184922
20. Khammash MH. Perfect adaptation in biology. *Cell Syst.* **2021**, *12*, 509–521. DOI:10.1016/j.cels.2021.05.020
21. López-Maury L, Marguerat S, Bähler J. Tuning gene expression to changing environments: From rapid responses to evolutionary adaptation. *Nat. Rev. Genet.* **2008**, *9*, 583–593. DOI:10.1038/nrg2398
22. El-Samad H, Goff JP, Khammash M. Calcium Homeostasis and Parturient Hypocalcemia: An Integral Feedback Perspective. *J. Theor. Biol.* **2002**, *214*, 17–29. DOI:10.1006/jtbi.2001.2422
23. Bhattacharya P, Raman K, Tangirala AK. On biological networks capable of robust adaptation in the presence of uncertainties: A linear systems-theoretic approach. *Math. Biosci.* **2023**, *358*, 108984. DOI:10.1016/j.mbs.2023.108984
24. Shi W, Ma W, Xiong L, Zhang M, Tang C. Adaptation with transcriptional regulation. *Sci. Rep.* **2017**, *7*, 42648. DOI:10.1038/srep42648
25. de Nadal E, Ammerer G, Posas F. Controlling gene expression in response to stress. *Nat. Rev. Genet.* **2011**, *12*, 833–845. DOI:10.1038/nrg3055
26. Li Y, Arce A, Lucci T, Rasmussen RA, Lucks JB. Dynamic RNA synthetic biology: New principles, practices and potential. *RNA Biol.* **2023**, *20*, 817–829. DOI:10.1080/15476286.2023.2269508
27. Waters LS, Storz G. Regulatory RNAs in Bacteria. *Cell* **2009**, *136*, 615–628. DOI:10.1016/j.cell.2009.01.043
28. Chappell J, Takahashi MK, Lucks JB. Creating small transcription activating RNAs. *Nat. Chem. Biol.* **2015**, *11*, 214–220. DOI:10.1038/nchembio.1737
29. Xu JZ, Zhang JL, Zhang WG. Antisense RNA: The new favorite in genetic research. *J. Zhejiang Univ. Sci. B* **2018**, *19*, 739–749. DOI:10.1631/jzus.B1700594
30. Sczakiel G. The Design of Antisense RNA. *Antisense Nucleic Acid. Drug Dev.* **1997**, *7*, 439–444. DOI:10.1089/oli.1.1997.7.439
31. Hoynes-O'Connor A, Moon TS. Development of Design Rules for Reliable Antisense RNA Behavior in *E. coli*. *ACS Synth. Biol.* **2016**, *5*, 1441–1454. DOI:10.1021/acssynbio.6b00036
32. Chappell J, Watters KE, Takahashi MK, Lucks JB. A renaissance in RNA synthetic biology: New mechanisms, applications and tools for the future. *Curr. Opin. Chem. Biol.* **2015**, *28*, 47–56. DOI:10.1016/j.cbpa.2015.05.018
33. Yoo SM, Na D, Lee SY. Design and use of synthetic regulatory small RNAs to control gene expression in *Escherichia coli*. *Nat. Protoc.* **2013**, *8*, 1694–1707. DOI:10.1038/nprot.2013.105
34. Chappell J, Westbrook A, Verosloff M, Lucks JB. Computational design of small transcription activating RNAs for versatile and dynamic gene regulation. *Nat. Commun.* **2017**, *8*, 1051. DOI:10.1038/s41467-017-01082-6
35. Lee YJ, Kim SJ, Moon TS. Multilevel Regulation of Bacterial Gene Expression with the Combined STAR and Antisense RNA System. *ACS Synth. Biol.* **2018**, *7*, 853–865. DOI:10.1021/acssynbio.7b00322
36. McSweeney MA, Zhang Y, Styczynski MP. Short Activators and Repressors of RNA Toehold Switches. *ACS Synth. Biol.* **2023**, *12*, 681–688. DOI:10.1021/acssynbio.2c00641
37. Green AA, Silver PA, Collins JJ, Yin P. Toehold Switches: De-Novo-Designed Regulators of Gene Expression. *Cell* **2014**, *159*, 925–939. DOI:10.1016/j.cell.2014.10.002
38. Isaacs FJ, Dwyer DJ, Ding C, Pervouchine DD, Cantor CR, Collins JJ. Engineered riboregulators enable post-transcriptional control of gene expression. *Nat. Biotechnol.* **2004**, *22*, 841–847. DOI:10.1038/nbt986
39. Mutalik VK, Qi L, Guimaraes JC, Lucks JB, Arkin AP. Rationally designed families of orthogonal RNA regulators of translation. *Nat. Chem. Biol.* **2012**, *8*, 447–454. DOI:10.1038/nchembio.919
40. Ekdahl AM, Rojano-Nisimura AM, Contreras LM. Engineering Toehold-Mediated Switches for Native RNA Detection and Regulation in Bacteria. *J. Mol. Biol.* **2022**, *434*, 167689. DOI:10.1016/j.jmb.2022.167689
41. Tabuchi T, Yokobayashi Y. Cell-free riboswitches. *RSC Chem. Biol.* **2021**, *2*, 1430–1440. DOI:10.1039/D1CB00138H
42. Wang T, Simmel FC. Riboswitch-inspired toehold riboregulators for gene regulation in *Escherichia coli*. *Nucleic Acids Res.* **2022**, *50*, 4784–4798. DOI:10.1093/nar/gkac275
43. Lee PW, Maerkl SJ. Regulatory Components for Bacterial Cell-Free Systems Engineering. *ACS Synth. Biol.* **2024**, *13*, 3827–3841. DOI:10.1021/acssynbio.4c00574
44. Lucks JB, Qi L, Mutalik VK, Wang D, Arkin AP. Versatile RNA-sensing transcriptional regulators for engineering genetic networks. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 8617–8622. DOI:10.1073/pnas.1015741108
45. Simmons TR, Partipilo G, Buchser R, Stankes AC, Srivastava R, Chiu D, et al. Rewiring native post-transcriptional global regulators to achieve designer, multi-layered genetic circuits. *Nat. Commun.* **2024**, *15*, 8752. DOI:10.1038/s41467-024-52976-1
46. Pham A, Bassett S, Chen W, Da Silva NA. Assembly of Metabolons in Yeast Using Cas6-Mediated RNA Scaffolding. *ACS Synth. Biol.* **2023**, *12*, 1164–1174. DOI:10.1021/acssynbio.2c00650

47. Patil SD, Sharma R, Srivastava S, Navani NK, Pathania R. Downregulation of *yidC* in *Escherichia coli* by Antisense RNA Expression Results in Sensitization to Antibacterial Essential Oils Eugenol and Carvacrol. *PLoS ONE* **2013**, *8*, e57370. DOI:10.1371/journal.pone.0057370
48. Barrangou R, Marraffini LA. CRISPR-Cas Systems: Prokaryotes Upgrade to Adaptive Immunity. *Mol. Cell* **2014**, *54*, 234–244. DOI:10.1016/j.molcel.2014.03.011
49. Qi LS, Larson MH, Gilbert LA, Doudna JA, Weissman JS, Arkin AP, et al. Repurposing CRISPR as an RNA-Guided Platform for Sequence-Specific Control of Gene Expression. *Cell* **2013**, *152*, 1173–1183. DOI:10.1016/j.cell.2013.02.022
50. Gilbert LA, Larson MH, Morsut L, Liu Z, Brar GA, Torres SE, et al. CRISPR-Mediated Modular RNA-Guided Regulation of Transcription in Eukaryotes. *Cell* **2013**, *154*, 442–451. DOI:10.1016/j.cell.2013.06.044
51. Nielsen AA, Voigt CA. Multi-input CRISPR/Cas genetic circuits that interface host regulatory networks. *Mol. Syst. Biol.* **2014**, *10*, 11:763. DOI:10.15252/msb.20145735
52. Zhao M, Kim J, Jiao J, Lim Y, Shi X, Guo S, et al. Construction of multilayered gene circuits using de-novo-designed synthetic transcriptional regulators in cell-free systems. *J. Biol. Eng.* **2024**, *18*, 64. DOI:10.1186/s13036-024-00459-8
53. Mitkas AA, Valverde M, Chen W. Dynamic modulation of enzyme activity by synthetic CRISPR–Cas6 endonucleases. *Nat. Chem. Biol.* **2022**, *18*, 492–500. DOI:10.1038/s41589-022-01005-7
54. Xi C, Chiu S, Voje WE, Carothers JM, Moon TS. Conditional guide RNA deactivation by mRNA and small molecule triggers in *Saccharomyces cerevisiae*. *New Biotechnol.* **2025**, *89*, 105–118. DOI:10.1016/j.nbt.2025.07.004
55. Westbrook A, Tang X, Marshall R, Maxwell CS, Chappell J, Agrawal DK, et al. Distinct timescales of RNA regulators enable the construction of a genetic pulse generator. *Biotech. Bioeng.* **2019**, *116*, 1139–1151. DOI:10.1002/bit.26918
56. Cameron DE, Bashor CJ, Collins JJ. A brief history of synthetic biology. *Nat. Rev. Microbiol.* **2014**, *12*, 381–390. DOI:10.1038/nrmicro3239
57. Elowitz MB, Leibler S. A synthetic oscillatory network of transcriptional regulators. *Nature* **2000**, *403*, 335–338. DOI:10.1038/35002125
58. Brophy JAN, Voigt CA. Principles of genetic circuit design. *Nat. Methods* **2014**, *11*, 508–520. DOI:10.1038/nmeth.2926
59. Khalil AS, Lu TK, Bashor CJ, Ramirez CL, Pyenson NC, Joung JK, et al. A Synthetic Biology Framework for Programming Eukaryotic Transcription Functions. *Cell* **2012**, *150*, 647–658. DOI:10.1016/j.cell.2012.05.045
60. Garg A, Lohmueller JJ, Silver PA, Armel TZ. Engineering synthetic TAL effectors with orthogonal target sites. *Nucleic Acids Res.* **2012**, *40*, 7584–7595. DOI:10.1093/nar/gks404
61. Lohmueller JJ, Armel TZ, Silver PA. A tunable zinc finger-based framework for Boolean logic computation in mammalian cells. *Nucleic Acids Res.* **2012**, *40*, 5180–5187. DOI:10.1093/nar/gks142
62. Li HS, Israni DV, Gagnon KA, Gan KA, Raymond MH, Sander JD, et al. Multidimensional control of therapeutic human cell function with synthetic gene circuits. *Science* **2022**, *378*, 1227–1234. DOI:10.1126/science.ade0156
63. Rhodius VA, Segall-Shapiro TH, Sharon BD, Ghodasara A, Orlova E, Tabakh H, et al. Design of orthogonal genetic switches based on a crosstalk map of σ s, anti- σ s, and promoters. *Mol. Syst. Biol.* **2013**, *9*, 702. DOI:10.1038/msb.2013.58
64. Shis DL, Bennett MR. Library of synthetic transcriptional AND gates built with split T7 RNA polymerase mutants. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 5028–5033. DOI:10.1073/pnas.1220157110
65. Segall-Shapiro TH, Meyer AJ, Ellington AD, Sontag ED, Voigt CA. A ‘resource allocator’ for transcription based on a highly fragmented T7 RNA polymerase. *Mol. Syst. Biol.* **2014**, *10*, 7:742. DOI:10.15252/msb.20145299
66. Kar S, Gardner EC, Javanmardi K, Boutz DR, Shroff R, Horton AP, et al. Directed evolution of an orthogonal transcription engine for programmable gene expression in eukaryotes. *iScience* **2025**, *28*, 111541. DOI:10.1016/j.isci.2024.111541
67. Grilly C, Stricker J, Pang WL, Bennett MR, Hasty J. A synthetic gene network for tuning protein degradation in *Saccharomyces cerevisiae*. *Mol. Syst. Biol.* **2007**, *3*, 127. DOI:10.1038/msb4100168
68. Cameron DE, Collins JJ. Tunable protein degradation in bacteria. *Nat. Biotechnol.* **2014**, *32*, 1276–1281. DOI:10.1038/nbt.3053
69. Andersen JB, Sternberg C, Poulsen LK, Bjørn SP, Givskov M, Molin S. New Unstable Variants of Green Fluorescent Protein for Studies of Transient Gene Expression in Bacteria. *Appl. Environ. Microbiol.* **1998**, *64*, 2240–2246. DOI:10.1128/AEM.64.6.2240-2246.1998
70. McGinness KE, Baker TA, Sauer RT. Engineering Controllable Protein Degradation. *Mol. Cell* **2006**, *22*, 701–707. DOI:10.1016/j.molcel.2006.04.027
71. Tague N, Coriano-Ortiz C, Sheets MB, Dunlop MJ. Light-inducible protein degradation in *E. coli* with the LOVdeg tag. *eLife* **2024**, *12*, RP87303. DOI:10.7554/eLife.87303
72. Verbič A, Lebar T, Praznik A, Jerala R. Subunits of an E3 Ligase Complex as Degrons for Efficient Degradation of Cytosolic, Nuclear, and Membrane Proteins. *ACS Synth. Biol.* **2024**, *13*, 792–803. DOI:10.1021/acssynbio.3c00588

73. Bleris L, Xie Z, Glass D, Adadey A, Sontag E, Benenson Y. Synthetic incoherent feedforward circuits show adaptation to the amount of their genetic template. *Mol. Syst. Biol.* **2011**, *7*, 519. DOI:10.1038/msb.2011.49
74. Guo S, Murray RM. Construction of Incoherent Feedforward Loop Circuits in a Cell-Free System and in Cells. *ACS Synth. Biol.* **2019**, *8*, 606–610. DOI:10.1021/acssynbio.8b00493
75. Martella A, Firth M, Taylor BJM, Göppert A, Cuomo EM, Roth RG, et al. Systematic Evaluation of CRISPRa and CRISPRi Modalities Enables Development of a Multiplexed, Orthogonal Gene Activation and Repression System. *ACS Synth. Biol.* **2019**, *8*, 1998–2006. DOI:10.1021/acssynbio.8b00527
76. McCarty NS, Graham AE, Studená L, Ledesma-Amaro R. Multiplexed CRISPR technologies for gene editing and transcriptional regulation. *Nat. Commun.* **2020**, *11*, 1281. DOI:10.1038/s41467-020-15053-x
77. Pacalin NM, Steinhart Z, Shi Q, Belk JA, Dorovskiy D, Kraft K, et al. Bidirectional epigenetic editing reveals hierarchies in gene regulation. *Nat. Biotechnol.* **2025**, *43*, 355–368. DOI:10.1038/s41587-024-02213-3
78. Kim SK, Woo SG, Park JH, Lee SG, Lee DH. Modular and signal-responsive transcriptional regulation using CRISPRi-aided genetic switches in *Escherichia coli*. *J. Biol. Eng.* **2025**, *19*, 56. DOI:10.1186/s13036-025-00526-8
79. Lian J, Hamedirad M, Hu S, Zhao H. Combinatorial metabolic engineering using an orthogonal tri-functional CRISPR system. *Nat. Commun.* **2017**, *8*, 1688. DOI:10.1038/s41467-017-01695-x
80. Chen Y, Niu X, Cheng M, Wang L, Sun P, Song H, et al. CRISPR/dCas9-RpoD-Mediated Simultaneous Transcriptional Activation and Repression in *Shewanella oneidensis* MR-1. *ACS Synth. Biol.* **2022**, *11*, 2184–2192. DOI:10.1021/acssynbio.2c00149
81. Chen Y, Li M, Liu X, Duan Q, Xiao L, Wang L, et al. Establishment of CRISPR-STAR System to Realise Simultaneous Transcriptional Activation and Repression in *Yarrowia lipolytica*. *Microb. Biotechnol.* **2025**, *18*, e70151. DOI:10.1111/1751-7915.70151
82. Huang HH, Bellato M, Qian Y, Cárdenas P, Pasotti L, Magni P, et al. dCas9 regulator to neutralize competition in CRISPRi circuits. *Nat. Commun.* **2021**, *12*, 1692. DOI:10.1038/s41467-021-21772-6
83. Elowitz MB, Levine AJ, Siggia ED, Swain PS. Stochastic Gene Expression in a Single Cell. *Science* **2002**, *297*, 1183–1186. DOI:10.1126/science.1070919
84. Paulsson J. Summing up the noise in gene networks. *Nature* **2004**, *427*, 415–418. DOI:10.1038/nature02257
85. Bashor CJ, Collins JJ. Understanding Biological Regulation Through Synthetic Biology. *Annu. Rev. Biophys.* **2018**, *47*, 399–423. DOI:10.1146/annurev-biophys-070816-033903
86. Sechkar K, Steel H. Model-guided gene circuit design for engineering genetically stable cell populations in diverse applications. *J. R. Soc. Interface.* **2025**, *22*, 20240602. DOI:10.1098/rsif.2024.0602
87. Del Vecchio D, Dy AJ, Qian Y. Control theory meets synthetic biology. *J. R. Soc. Interface.* **2016**, *13*, 20160380. DOI:10.1098/rsif.2016.0380
88. Sechkar K, Steel H. Combining positive and negative regulation for modular and robust biomolecular control architectures. *bioRxiv* **2024**, 2024-03. DOI:10.1101/2024.03.22.586143
89. Hu CY, Varner JD, Lucks JB. Generating Effective Models and Parameters for RNA Genetic Circuits. *ACS Synth. Biol.* **2015**, *4*, 914–926. DOI:10.1021/acssynbio.5b00077
90. Gillespie DT. Exact stochastic simulation of coupled chemical reactions. *J. Phys. Chem.* **1977**, *81*, 2340–2361. DOI:10.1021/j100540a008
91. Raj A, van Oudenaarden A. Nature, Nurture, or Chance: Stochastic Gene Expression and Its Consequences. *Cell* **2008**, *135*, 216–226. DOI:10.1016/j.cell.2008.09.050
92. Fang Z, Gupta A, Kumar S, Khammash M. Advanced methods for gene network identification and noise decomposition from single-cell data. *Nat. Commun.* **2024**, *15*, 4911. DOI:10.1038/s41467-024-49177-1
93. Goetz H, Zhang R, Wang X, Tian XJ. Resource competition-driven bistability and stochastic switching amplify gene expression noise. *PLoS Comput. Biol.* **2025**, *21*, e1012931. DOI:10.1371/journal.pcbi.1012931
94. Schnoerr D, Sanguinetti G, Grima R. Approximation and Inference Methods for Stochastic Biochemical Kinetics. *J. Phys. A Math. Theor.* **2017**, *50*, 093001. DOI:10.1088/1751-8121/aab38b
95. Thattai M, van Oudenaarden A. Intrinsic noise in gene regulatory networks. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 8614–8619. DOI:10.1073/pnas.151588598
96. Chu BK, Tse MJ, Sato RR, Read EL. Markov State Models of gene regulatory networks. *BMC Syst. Biol.* **2017**, *11*, 14. DOI:10.1186/s12918-017-0394-4
97. Swain PS, Elowitz MB, Siggia ED. Intrinsic and extrinsic contributions to stochasticity in gene expression. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 12795–12800. DOI:10.1073/pnas.162041399
98. Santos-Moreno J, Tasiudi E, Stelling J, Schaerli Y. Multistable and dynamic CRISPRi-based synthetic circuits. *Nat. Commun.* **2020**, *11*, 2746. DOI:10.1038/s41467-020-16574-1

99. Briat C, Gupta A, Khammash M. Antithetic Integral Feedback Ensures Robust Perfect Adaptation in Noisy Biomolecular Networks. *Cell Syst.* **2016**, 2, 15–26. DOI:10.1016/j.cels.2016.01.004
100. Salis HM. The Ribosome Binding Site Calculator. In *Methods in Enzymology*; Academic Press Inc.: Cambridge, MA, USA, 2011; Volume 498, pp. 19–42.
101. McNerney MP, Zhang Y, Steppe P, Silverman AD, Jewett MC, Styczynski MP. Point-of-care biomarker quantification enabled by sample-specific calibration. *Sci. Adv.* **2019**, 5, eaax4473. DOI:10.1126/sciadv.aax4473
102. Jayaraman P, Devarajan K, Chua TK, Zhang H, Gunawan E, Poh CL. Blue light-mediated transcriptional activation and repression of gene expression in bacteria. *Nucleic Acids Res.* **2016**, 44, 6994–7005. DOI:10.1093/nar/gkw548
103. Santos-Moreno J, Schaerli Y. CRISPR-based gene expression control for synthetic gene circuits. *Biochem. Soc. Trans.* **2020**, 48, 1979–1993. DOI:10.1042/BST20200020
104. Pandi A, Koch M, Voyvodic PL, Soudier P, Bonnet J, Kushwaha M, et al. Metabolic perceptrons for neural computing in biological systems. *Nat. Commun.* **2019**, 10, 3880. DOI:10.1038/s41467-019-11889-0
105. Moghimianavval H, Gispert I, Castillo SR, Corning OBWH, Liu AP, Cuba Samaniego C. Engineering Sequestration-Based Biomolecular Classifiers with Shared Resources. *ACS Synth. Biol.* **2024**, 13, 3231–3245. DOI:10.1021/acssynbio.4c00270
106. Cherry KM, Qian L. Scaling up molecular pattern recognition with DNA-based winner-take-all neural networks. *Nature* **2018**, 559, 370–376. DOI:10.1038/s41586-018-0289-6
107. Rizik L, Danial L, Habib M, Weiss R, Daniel R. Synthetic neuromorphic computing in living cells. *Nat. Commun.* **2022**, 13, 5602. DOI:10.1038/s41467-022-33288-8