

Article

# Production of Highly Modified C<sub>30</sub>-carotenoids with Singlet Oxygen-quenching Activities, 5-glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-oic Acid, and Its Three Intermediates Using Genes from *Planococcus maritimus* Strain iso-3

Moe Hagiwara <sup>1</sup>, Chinatsu Maehara <sup>1</sup>, Miho Takemura <sup>2</sup>, Norihiko Misawa <sup>2,\*</sup>, and Kazutoshi Shindo <sup>1,\*</sup>

<sup>1</sup> Department of Food and Nutrition, Japan Women's University, 2-8-1 Mejirodai, Bunkyo-ku, Tokyo 112-8681, Japan; m1702023hm@ug.jwu.ac.jp (M.H.); m1803045mc@ug.jwu.ac.jp (C.M.)

<sup>2</sup> Research Institute for Bioresources and Biotechnology, Ishikawa Prefectural University, 1-308 Suematsu, Nonoichi-shi, Ishikawa 921-8836, Japan; mtake@ishikawa-pu.ac.jp (M.T.)

\* Corresponding authors. E-mail: n-misawa@ishikawa-pu.ac.jp (N.M.); kshindo@fc.jwu.ac.jp (K.S.)

Received: 12 December 2022; Accepted: 3 February 2023; Available online: 7 February 2023

**ABSTRACT:** *Planococcus maritimus* strain iso-3 was previously isolated from intertidal sediment in the North Sea and was found to produce a highly modified C<sub>30</sub>-carotenoid, methyl-5-glucosyl-5,6-dihydro-4,4'-diapolycopenoate, as the final product. In this study, we analyzed the function of the carotenoid terminal oxidase *crtP* (renamed *cruO*) and aldehyde dehydrogenase *aldH* genes in *P. maritimus* strain iso-3 and elucidated the carotenoid biosynthetic pathway for this strain at the gene level. We produced four novel C<sub>30</sub>-carotenoids with potent singlet oxygen-quenching activities, 5-glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-oic acid and its three intermediates, which were obtained using *E. coli* cells carrying the *cruO* (and *aldH*) gene(s) in addition to the known *P. maritimus* carotenogenic genes.

**Keywords:** C<sub>30</sub>-carotenoids; Marine bacterium; *Planococcus*; Diapolycopenoic acid; Singlet oxygen-quenching activity




© 2023 The authors. This is an open access article under the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

C<sub>30</sub>-Carotenoids are acyclic isoprenoid pigments composed of 30 carbons as the basic backbone structure and have been found only in non-photosynthetic bacteria [1]. The biosynthetically first yellow-colored C<sub>30</sub>-carotenoid 4,4'-diaponeurosporene is synthesized from farnesyl diphosphate (FPP) by way of 15-*cis*-4,4'-diapophytoene with the action of 4,4'-diapophytoene synthase (CrtM) and 4,4'-diapophytoene desaturase [CrtN (CrtNa)] (Figure 1, [2,3]). This biosynthetic pathway is common in all C<sub>30</sub>-carotenogenic bacteria.

*Planococcus maritimus* strain iso-3, a yellow marine bacterium belonging to the Firmicutes family, was isolated from intertidal sediment in the North Sea, which is industrially polluted [4]. This bacterium was shown to produce a highly modified C<sub>30</sub>-carotenoid, methyl 5-glucosyl-5,6-dihydro-4,4'-diapolycopenoate, as the final product [5]. We analyzed the structure and function of its carotenoid biosynthesis gene cluster through structural determination of carotenoids that were produced in *Escherichia coli* expressing combinations of the gene candidates found in the gene cluster [3]. It was consequently shown that 4,4'-diaponeurosporene was converted to 5-hydroxy-5,6-dihydro-4,4'-diaponeurosporene with carotenoid hydratase (CruF) that shared significant homology to the *Deinococcus* CruF [6], and then metabolized to 5-hydroxy-5,6-dihydro-4,4'-diapolycopenoate with 5-hydroxy-5,6-dihydro-4,4'-diaponeurosporene desaturase (CrtNb) (Figure 1). The glucosyl transferase (*GT*) [7–9] has also been found in the *Planococcus* carotenoid gene cluster [3].



**Figure 1.** Elucidated carotenoid biosynthetic pathway of *Planococcus maritimus* strain iso-3. Only an *MT* gene is not obtained.

In this study, we analyzed the structure and function of the carotenoid oxidase *crtP* (renamed *cruO*) and aldehyde dehydrogenase *aldH* genes, which were newly isolated from places different from the above-mentioned gene cluster in *P. maritimus* iso-3 and elucidated the carotenoid biosynthetic pathway for 5-glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-oic acid at the gene level. Additionally, we demonstrated four novel C<sub>30</sub>-carotenoids that were produced in *E. coli* expressing combinations of *P. maritimus* carotenoid biosynthesis genes with their singlet oxygen-quenching activities.

## 2. Materials and Methods

### 2.1. Bacterial Strains and Growth Conditions

*E. coli* K12 DH5 $\alpha$  and JM101(DE3) cells were used for DNA manipulation and expression of carotenoid biosynthesis genes, respectively. These *E. coli* strains and their transformants were grown in 2 × YT (2YT) medium (16 g/L of tryptone, 10 g/L of yeast extract, 5 g/L of NaCl) containing 10 mg/L of tetracycline (as needed) at 37 °C or 21 °C.

## 2.2. DNA Isolation of *Planococcus maritimus* Strain iso-3

Genomic DNA was prepared from *P. maritimus* strain iso-3 according to the method described by Nishida et al. [10].

## 2.3. Cloning of the *crtP* (*cruO*) and *aldH* Genes from *P. maritimus* Strain iso-3


Based on the sequences of the *crtP* and *aldH* genes in the genomes of *Planococcus faecalis* and *Planococcus plakortidis*, primers containing the restriction sites were designed, as shown in Table S1. The coding regions of individual genes were amplified by PCR (polymerase chain reaction) using genomic DNAs. PCR products were cloned into a plasmid vector and sequenced as described [3].

## 2.4. Sequence Analysis

Homology search was performed using BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). Amino acid alignment was performed using MAFFT (<http://www.mafft.cbrc.jp/>) and phylogenetic tree was constructed using the neighbor joining method, both as implemented on the GenomeNet (<https://www.genome.jp/tools/ete/>).

## 2.5. Expression of the *crtP* (*cruO*) and *aldH* Genes from *P. maritimus* Strain iso-3 in *E. coli*

The coding region of the *crtP* (*cruO*) gene from *P. maritimus* strain iso-3 was inserted into the plasmids pAC-HIMNFnB and pAC-HIMNFnB-G [3]. These plasmids were designated pAC-HIMNFnBP and pAC-HIMNFnBGP, respectively (Figure 2). The coding region of the *aldH* gene from *P. maritimus* strain iso-3 was independently inserted into pAC-HIMNFnBP and pAC-HIMNFnBGP. The resultant plasmids were named pAC-HIMNFnBPA and pAC-HIMNFnB-GPA (Figure 2). All plasmids were introduced into wild-type *E. coli* (JM101 (DE3)). The transformed *E. coli* strains were grown in 2YT medium at 37 °C. The following day, the culture was inoculated in a new 2YT medium (100 ml medium in a 500 ml Sakaguchi flask) and cultured at 21 °C for two days.



**Figure 2.** Structure of plasmids constructed in this study.

## 2.6. Extraction and High-Performance Liquid Chromatography (HPLC) Analysis of Carotenoids from *E. coli* Cells

Extraction of carotenoids from recombinant *E. coli* was performed as described by Fraser et al. [11]. *E. coli* cultures were centrifuged ( $8000g \times 5$  min). Cell pellets were extracted with methanol (MeOH) using a mixer for 5 min. Tris-HCl (50 mM, pH 7.5) and 1 M NaCl were added and mixed. Chloroform was added to the mixture and mixed for 5 min. After centrifugation, the chloroform phase was collected and dried through centrifugal evaporation. Dried residues were resuspended in ethyl acetate (EtOAc) and subjected to HPLC with a Waters Alliance 2695-2996 (PDA) system (Waters, Milford, MA, USA). HPLC was performed according to the method described by Yokoyama and Miki [12] using TSKgel ODS-80Ts ( $4.6 \times 150$  mm,  $5 \mu\text{m}$ ; Tosoh, Tokyo, Japan). Briefly, the extract was eluted at a flow rate of 1.0 mL/min at 25 °C with solvent A [water ( $\text{H}_2\text{O}$ )-MeOH, 5:95] for 5 min, followed by a linear gradient from solvent A to solvent B (tetrahydrofuran-MeOH, 3:7) for 5 min and solvent B alone for 8 min. The produced carotenoids were identified by comparing their retention times and absorption spectra with those of our authentic standards. When the produced carotenoids were not compounds in our authentic standards, as described in the following sections, we isolated the produced carotenoids and determined their structures using high-resolution electrospray ionization mass spectrometry (HRESI-MS) and nuclear magnetic resonance (NMR) analyses.

## 2.7. Isolation of Respective Highly Modified $C_{30}$ -Carotenoids

### 2.7.1. 5-Hydroxy-5,6-dihydro-4,4'-diapolycopen-4'-al (1)

The transformed *E. coli* cells carrying pAC-HIMNFNBp were collected using centrifugation from 2 L culture and extracted with 540 mL acetone-MeOH (7:2) and 600 mL dichloromethane ( $\text{CH}_2\text{Cl}_2$ )-MeOH (1:1) with sonication in a stepwise manner. The combined extract (1140 mL) was concentrated to a small volume *in vacuo* and partitioned with EtOAc/ $\text{H}_2\text{O}$  (300 mL/150 mL). The EtOAc layer containing the produced carotenoids was evaporated to dryness to obtain yellow oil (125.0 mg). *n*-Hexane (3 mL) was added to the oil and sonicated for 3 min to remove fats and fatty acids ( $\times 3$ ). The precipitate (10.1 mg) was subjected to preparative ODS HPLC [column: Develosil C30-UG (20 mm  $\times$  250 mm, Nomura Chemical, Co. Ltd, Aichi, Japan), solvent: acetonitrile ( $\text{CH}_3\text{CN}$ )-tetrahydrofuran (THF) (3:2), flow rate: 8.0 mL/min, detection: PDA (250–700 nm)]. The peak at  $t_R$  9.6 min was collected and concentrated to dryness to afford pure 5-hydroxy-5,6-dihydro-4,4'-diapolycopen-4'-al (1) (1.1 mg).

### 2.7.2. 5-Hydroxy-5,6-dihydro-4,4'-diapolycopen-4'-oic Acid (2)

Transformed *E. coli* cells carrying pAC-HIMNFNBPA were collected by centrifugation from 2 L of culture and extracted with acetone-MeOH (7:2) and  $\text{CH}_2\text{Cl}_2$ -MeOH (1:1) by stepwise sonication. The extract (1140 mL) was concentrated to a small volume *in vacuo* and partitioned with EtOAc/ $\text{H}_2\text{O}$  (300 mL/150 mL). The EtOAc layer was evaporated to dryness to obtain a yellow oil (113.6 mg). *n*-Hexane (3 mL) was added to the oil and sonicated for 3 min to remove fats and fatty acids ( $\times 3$ ). The precipitate (17.9 mg) was subjected to preparative ODS HPLC [column: Develosil C30-UG (20 mm  $\times$  250 mm), solvent: 90% (*v/v*)  $\text{CH}_3\text{CN}$  containing 20 mM  $\text{H}_3\text{PO}_4$ -THF (3:2), flow rate: 8.0 mL/min, detection: PDA (250–700 nm)]. The peak at  $t_R$  9.8 min was collected and concentrated to 5 mL and partitioned between EtOAc/ $\text{H}_2\text{O}$  (100 mL each). The EtOAc layer was concentrated to dryness to obtain pure 5-hydroxy-5,6-dihydro-4,4'-diapolycopen-4'-oic acid (2) (1.8 mg).

### 2.7.3. 5-Glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-al (3)

The transformed *E. coli* cells carrying pAC-HIMNFNBGP were collected using centrifugation from 2 L of culture and extracted with 540 mL acetone-MeOH (7:2) and 600 mL  $\text{CH}_2\text{Cl}_2$ -MeOH (1:1) with sonication in a stepwise manner. The combined extract (1140 mL) was concentrated to a small volume *in vacuo* and partitioned with EtOAc/ $\text{H}_2\text{O}$  (300 mL/150 mL). The EtOAc layer containing the produced carotenoids was evaporated to dryness to obtain a yellow oil (301.7 mg). *n*-Hexane (3 mL) was added to the oil and sonicated for 3 min to remove fatty acids ( $\times 3$ ). The precipitate (142.2 mg) was subjected to preparative ODS HPLC [column: Develosil C30-UG (20 mm  $\times$  250 mm), solvent:  $\text{CH}_3\text{CN}$ -THF (3:2), flow rate: 8.0 mL/min, detection: PDA (250–700 nm)]. The peak at  $t_R$  8.2 min was collected and concentrated to dryness to afford pure 5-glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-al (3) (1.2 mg).

### 2.7.4. 5-Glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-oic Acid (4)

Transformed *E. coli* cells carrying pAC-HIMNFNBGPA were collected using centrifugation from 2 L of culture and extracted with acetone-MeOH (7:2) and  $\text{CH}_2\text{Cl}_2$ -MeOH (1:1) using stepwise sonication. The extract (1140 mL) was concentrated to a small volume *in vacuo* and partitioned with EtOAc/ $\text{H}_2\text{O}$  (300 mL/150 mL). The ethyl acetate (EtOAc) layer was evaporated to dryness (203.2 mg). *n*-Hexane (3 mL) was added to the oil and sonicated for 3 min to remove fats and fatty acids ( $\times 3$ ). The precipitate (58.9 mg) was subjected to preparative ODS HPLC [column: Develosil C30-UG (20 mm  $\times$  250 mm), solvent: 90% (*v/v*)  $\text{CH}_3\text{CN}$  containing 20 mM  $\text{H}_3\text{PO}_4$ -THF (3:2), flow rate: 8.0 mL/min, detection: PDA (250–700 nm)]. The peak at  $t_R$  11.2 min was collected

and concentrated to 5 mL and partitioned between EtOAc/H<sub>2</sub>O (each 100 mL). The EtOAc layer was concentrated to dryness to obtain pure 5-glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-oic acid (**4**) (3.2 mg).

## 2.8. Singlet Oxygen-Quenching Activity

For the measurement of singlet oxygen-quenching activity, 80  $\mu$ L of 25  $\mu$ M methylene blue and 100  $\mu$ L of 0.24 M linoleic acid, with or without 40  $\mu$ L of carotenoid (final concentration, 1–25  $\mu$ M; each dissolved in ethanol), were added to 5 mL glass test tubes. The tubes were mixed well and illuminated at 7000 lx and 22 °C for 3 h in a Styrofoam box. Subsequently, 120  $\mu$ L of the reaction mixture was removed and diluted to 3.48 mL with ethanol. The OD<sub>235</sub> was measured to estimate the formation of conjugated dienes. OD<sub>235</sub> in the absence of carotenoids was measured as a negative control [no singlet oxygen (<sup>1</sup>O<sub>2</sub>)-quenching activity]. The <sup>1</sup>O<sub>2</sub>-quenching activity of carotenoids was calculated from OD<sub>235</sub> in the presence of carotenoids relative to this reference value.

The activity is indicated as the IC<sub>50</sub> value, which represents the concentration at which 50% inhibition occurs.

## 2.9. Physico-Chemical Properties of Respective Highly Modified C<sub>30</sub>-Carotenoids

### 2.9.1. 5-Hydroxy-5,6-dihydro-4,4'-diapolycopen-4'-al (**1**)

HR-ESI-MS (+) *m/z* 455.29275 (C<sub>30</sub>H<sub>40</sub>O<sub>2</sub>Na; cald. for 455.29260,  $\Delta$  0.33 ppm). UV-Vis  $\lambda_{\text{max}}$  ( $\epsilon$ ) in MeOH 254 (9200), 315 (6800), 407 (43000), 428 (62000), 452 (56000). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$ : 1.14 (3H, s, H-4), 1.14 (3H, s, H-18), 1.90 (3H, s, H-18'), 1.94 (3H, s, H-19), 1.98 (3H, s, H-20), 1.98 (3H, s, H-20'), 2.02 (3H, s, H-19'), 2.32 (2H, d, *J* = 7.6 Hz, H-6), 5.77 (1H, dd, *J* = 7.6, 15.1 Hz, H-7), 6.14 (1H, d, *J* = 12.0 Hz, H-10), 6.22 (1H, d, *J* = 15.1 Hz, H-8), 6.26 (1H, d, *J* = 14.5 Hz, H-14), 6.33 (1H, d, *J* = 15.0 Hz, H-14'), 6.36 (1H, d, *J* = 14.0 Hz, H-12), 6.43 (1H, d, *J* = 12.3 Hz, H-10'), 6.50 (1H, d, *J* = 15.0 Hz, H-12'), 6.63 (1H, dd, *J* = 12.0, 14.0 Hz, H-11), 6.65 (1H, dd, *J* = 10.7, 14.8 Hz, H-7'), 6.65 (1H, dd, *J* = 12.5, 15.0 Hz, H-11'), 6.65 (2H, H-15 and H-15'), 6.75 (1H, d, *J* = 14.8 Hz, H-8'), 6.94 (1H, d, *J* = 10.7 Hz, H-6'), 9.45 (1H, s, H-4') (Figure S1). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$ : 12.8 (C-19), 12.8 (C-20), 12.8 (C-18'), 12.8 (C-20'), 13.0 (C-19'), 29.2 (C-4), 29.3 (C-18), 47.4 (C-6), 71.0 (C-5), 122.4 (C-7'), 124.4 (C-11'), 124.8 (C-11), 125.3 (C-7), 129.8 (C-15')<sup>c</sup>, 131.0 (C-10), 131.2 (C-15)<sup>c</sup>, 132.5 (C-14)<sup>b</sup>, 134.8 (C-14)<sup>b</sup>, 134.9 (C-9'), 135.4 (C-13)<sup>a</sup>, 136.1 (C-13)<sup>a</sup>, 136.6 (C-5'), 137.3 (C-9), 137.6 (C-12), 137.8 (C-10'), 138.7 (C-8), 141.0 (C-12'), 146.1 (C-8'), 149.4 (C-6'), 194.6 (C-4') (Figure S2). <sup>a, b, c</sup>: Interchangeable.

### 2.9.2. 5-Hydroxy-5,6-dihydro-4,4'-diapolycopen-4'-oic Acid (**2**)

HR-ESI-MS (+) *m/z* 471.28902 (C<sub>30</sub>H<sub>40</sub>O<sub>3</sub>Na; cald. for 471.28751,  $\Delta$  1.51 ppm). UV-Vis  $\lambda_{\text{max}}$  ( $\epsilon$ ) in MeOH 254 (9200), 315 (6800), 407 (43000), 428 (62000), 452 (56000). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>)  $\delta$ : 1.06 (3H, s, H-4), 1.06 (3H, s, H-18), 1.87 (3H, s, H-19), 1.88 (3H, s, H-18'), 1.93 (3H, s, H-20), 1.94 (3H, s, H-20'), 1.96 (3H, s, H-19'), 2.18 (2H, d, *J* = 7.1 Hz, H-6), 5.77 (1H, td, *J* = 7.1, 16.0 Hz, H-7), 6.12 (1H, d, *J* = 16.0 Hz, H-8), 6.13 (1H, d, *J* = 10.2 Hz, H-10), 6.32 (1H, d, *J* = 10.4 Hz, H-14), 6.36 (1H, d, *J* = 15.0 Hz, H-12), 6.38 (1H, d, *J* = 11.0 Hz, H-14'), 6.53 (1H, dd, *J* = 11.2, 15.3 Hz, H-7'), 6.40 (1H, d, *J* = 11.3 Hz, H-10'), 6.48 (1H, d, *J* = 14.8 Hz, H-12'), 6.66 (1H, dd, *J* = 10.2, 15.0 Hz, H-11), 6.68 (1H, dd, *J* = 11.3, 14.8 Hz, H-11'), 6.72 (1H, dd, *J* = 11.0, 15.0 Hz, H-15'), 6.73 (1H, dd, *J* = 10.4, 15.0 Hz, H-15), 7.12 (1H, d, *J* = 11.2 Hz, H-6') (Figure S3). <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>)  $\delta$ : 12.7 (C-20'), 12.8 (C-20), 12.8 (C-19'), 13.0 (C-19), 13.3 (C-18'), 29.4 (C-18), 29.4 (C-4), 47.6 (C-6), 69.5 (C-5), 124.1 (C-7'), 125.2 (C-11), 125.6 (C-11'), 127.2 (C-7), 128.6 (C-5'), 130.3 (C-10), 130.3 (C-15'), 131.2 (C-15), 132.6 (C-14), 134.0 (C-14'), 135.2 (C-10'), 135.6 (C-9), 135.6 (C-9'), 135.6 (C-13'), 136.8 (C-13), 136.8 (C-6'), 136.9 (C-8), 137.2 (C-12), 139.4 (C-12'), 142.5 (C-8'), 169.7 (C-4') (Figure S4).

### 2.9.3. 5-Glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-al (**3**)

HR-ESI-MS (+): *m/z* 617.34744 (C<sub>36</sub>H<sub>50</sub>O<sub>7</sub>Na; cald. for 617.34542,  $\Delta$  3.27 ppm). UV-Vis  $\lambda_{\text{max}}$  ( $\epsilon$ ) in MeOH 254 (9200), 315 (6800), 407 (43000), 428 (62000), 452 (56000). <sup>1</sup>H NMR (CD<sub>3</sub>OD)  $\delta$ : 1.26 (3H, s, H-4), 1.26 (3H, s, H-18), 1.87 (3H, s, H-18'), 1.92 (3H, s, H-19), 1.97 (3H, s, H-20), 1.99 (3H, s, H-20'), 2.03 (3H, s, H-19'), 2.42 (2H, d, *J* = 7.5 Hz, H-6), 3.15 (1H, dd, *J* = 7.6, 9.0 Hz, H-2''), 3.24 (1H, m, H-5''), 3.26 (1H, m, H-4''), 3.35 (1H, dd, *J* = 8.9, 9.0 Hz, H-3''), 3.65 (1H, m, H-6''a), 3.82 (1H, m, H-6''b), 4.51 (1H, d, *J* = 7.6 Hz, H-1''), 5.91 (1H, m, H-7), 6.11 (1H, d, *J* = 12.0 Hz, H-10), 6.18 (1H, d, *J* = 15.2 Hz, H-8), 6.26 (2H, H-14 and H-14'), 6.37 (1H, d, *J* = 15.0 Hz, H-12), 6.51 (1H, d, *J* = 12.3 Hz, H-10'), 6.56 (1H, d, *J* = 15.0 Hz, H-12'), 6.65 (2H, H-15 and H-15'), 6.68 (1H, dd, *J* = 12.0, 15.0 Hz, H-11), 6.72 (1H, dd, *J* = 12.3, 14.9 Hz, H-11'), 6.74 (1H, dd, *J* = 10.8, 14.9 Hz, H-7'), 6.88 (1H, d, *J* = 14.9 Hz, H-8'), 7.10 (1H, d, *J* = 10.8 Hz, H-6'), 9.39 (1H, s, H-4') (Figure S5). <sup>13</sup>C NMR (CD<sub>3</sub>OD)  $\delta$ : 8.0 (C-18'), 11.2 (C-20), 11.3 (C-19')<sup>c</sup>, 11.5 (C-20')<sup>c</sup>, 11.6 (C-19), 25.3 (C-18), 25.6 (C-4), 29.3 (C-18), 45.3 (C-6), 61.4 (C-6''), 70.4 (C-4''), 73.8 (C-2''), 76.2 (C-5''), 76.8 (C-3''), 77.8 (C-5), 97.3 (C-1''), 122.1 (C-7'), 123.0 (C-11'), 123.7 (C-11), 125.6 (C-7), 129.7 (C-15)<sup>b</sup>, 130.3 (C-10), 131.2

(C-15)<sup>b</sup>, 132.2 (C-14)<sup>a</sup>, 134.8 (C-9'), 134.9 (C-14')<sup>a</sup>, 135.4 (C-9), 135.8 (C-13'), 135.9 (C-13), 136.1 (C-5'), 137.0 (C-12), 137.4 (C-8), 137.9 (C-10'), 141.0 (C-12'), 146.7 (C-8'), 150.4 (C-6'), 195.1 (C-4') (Figure S6). <sup>a, b, c</sup>: Interchangeable.

#### 2.9.4. 5-Glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-oic Acid (4)

HR-ESI-MS (+): *m/z* 633.33807 (C<sub>36</sub>H<sub>50</sub>O<sub>8</sub>Na; cald. for 633.34034, Δ 3.58 ppm). UV-Vis λ<sub>max</sub> (ε) in MeOH 254 (9200), 315 (6800), 407 (43000), 428 (62000), 452 (56000). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ: 1.14 (6H, s, H-4 and H-18), 1.88 (3H, s, H-19), 1.90 (3H, s, H-18'), 1.93 (3H, s, H-20'), 1.94 (3H, s, H-20), 1.97 (3H, s, H-19'), 2.32 (2H, t, *J*=7.5 Hz, H-6), 2.89 (1H, dd, *J*=7.6, 8.4 Hz, H-2''), 3.02 (1H, dd, *J*=8.4, 8.4 Hz, H-4''), 3.07 (1H, dd, *J*=5.2, 8.4 Hz, H-5''), 3.14 (1H, dd, *J*=8.4, 8.4 Hz, H-3''), 3.39 (1H, dd, *J*=5.2, 11.0 Hz, H-6''b), 3.62 (1H, d, *J*=11.0, H-6''a), 4.32 (1H, d, *J*=7.6 Hz, H-1''), 5.88 (1H, td, *J*=7.5, 15.6 Hz, H-7), 6.13 (1H, d, *J*=12.0 Hz, H-10), 6.14 (1H, d, *J*=15.6 Hz, H-8), 6.33 (1H, d, *J*=10.5 Hz, H-14), 6.37 (1H, d, *J*=15.0 Hz, H-12), 6.42 (1H, d, *J*=10.5 Hz, H-14'), 6.43 (1H, d, *J*=10.8 Hz, H-10'), 6.50 (1H, d, *J*=15.4 Hz, H-12'), 6.55 (1H, dd, *J*=11.0, 15.0 Hz, H-7'), 6.65 (1H, dd, *J*=12.0, 15.0 Hz, H-11), 6.68 (1H, dd, *J*=12.2, 15.0 Hz, H-11'), 6.71 (1H, d, *J*=15.0 Hz, H-8'), 6.73 (2H, H-15 and H-15'), 7.18 (1H, d, *J*=11.0 Hz, H-6') (Figure S7). <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ: 12.8 (C-20), 12.8 (C-19'), 12.8 (C-20'), 13.0 (C-18'), 13.1 (C-19), 26.4 (C-4), 26.9 (C-18), 45.1 (C-6), 61.4 (C-6''), 70.4 (C-4''), 73.8 (C-2''), 76.7 (C-5''), 76.9 (C-5), 77.2 (C-3''), 97.4 (C-1''), 123.5 (C-7'), 125.2 (C-11'), 125.7 (C-11), 126.7 (C-5'), 127.1 (C-7), 130.3 (C-10), 130.3 (C-15), 130.3 (C-15'), 132.6 (C-14), 134.2 (C-13'), 135.5 (C-14'), 135.5 (C-9'), 135.8 (C-10'), 136.3 (C-9), 137.0 (C-8), 137.0 (C-13), 137.2 (C-12), 138.3 (C-6'), 139.8 (C-12'), 143.8 (C-8'), 169.5 (C-4') (Figure S8).

### 3. Results and Discussion


#### 3.1. Isolation of the *crtP* and *aldH* Genes from *P. maritimus* Strain iso-3

Previously, we isolated the carotenoid biosynthesis gene cluster from *P. maritimus* iso-3, which contains *crtN* (*crtNa*), *crtM*, *crtNb*, *GT*, and *cruF* [3]. However, these genes were not sufficient to synthesize the carotenoid of *P. maritimus*, such as methyl 5-glucosyl-5,6-dihydro-4,4'-diapolycopenoate. Recently, Lee et al. cloned *crtP* and *aldH* genes from *P. faecalis* as carotenogenic genes [13]. Since the genome sequences are similar between the *Planococcus* genera, we designed PCR primers for these genes based on the genome sequences of *P. faecalis* and *P. plakorditis* (Table S1). PCR cloning with these primers resulted in the isolation of two genes from *P. maritimus* iso-3.

In *Staphylococcus aureus*, Pelz et al. found the gene mediating the oxidation of the terminal methyl group in a C<sub>30</sub>-carotenoid and named it *crtP* [8]. The *crtP* designation was adopted for the ortholog in *P. faecalis* [13]. In contrast, the *crtP* gene has been widely designated as the cyanobacterial phytoene desaturase gene [14]. Because the same gene name for distinct carotenoid genes is confusing, we propose that *crtP* be renamed *cruO* for the carotenoid terminal oxidase gene.

#### 3.2. Sequence Analysis of the Carotenoid Biosynthesis Gene Candidates

Subsequently, we performed a sequence analysis of the *cruO* and *aldH* genes isolated from *P. maritimus* iso-3. The predicted amino acid sequence of the *cruO* gene product (CruO) showed 86% identity to those of the *P. faecalis* CrtP, while CruO was 29%, 25%, and 23 % identical to CrtNb, CrtNa, and Orf2 that were isolated from the same *P. maritimus* iso-3, respectively. The phylogenetic tree of *crtN*-homologous genes (Table 1) showed that the *P. maritimus* iso-3 *cruO* gene encoding carotenoid terminal oxidase, along with the other corresponding genes (*crtP*), fell into the *crtNb* clade. This clade included the *crtNb* gene encoding carotenoid desaturase from *P. maritimus* iso-3 at earlier branch point (Figure 3). On the other hand, the predicted amino acid sequences of AldH showed 79% identity to those of the *P. faecalis* AldH. These *aldH* genes were found to belong to another gene family different from the *crtNc* (*aldH*) clade.



**Figure 3.** Phylogenetic tree of *crtN*-homologous genes.

Amino acid alignment of the encoded proteins was performed using MAFFT (<http://www.mafft.cbrc.jp/>) and phylogenetic trees was constructed using the neighbor joining method, both as implemented on the GenomeNet (<https://www.genome.jp/tools/ete/>). Accession numbers of the sequences are listed in Table 1.


**Table 1.** Examples of *crtN*-homologous genes and their accession numbers.

Organism	Gene	Accession No.
<i>Staphylococcus aureus</i>	<i>crtP</i> ( <i>crtNb</i> )	WP_102782088.1
<i>Staphylococcus aureus</i>	<i>crtN</i> ( <i>crtNa</i> )	WP_057520765.1
<i>Staphylococcus aureus</i>	<i>crtNc</i>	WP_001084326.1
<i>Halobacillus halophilus</i>	<i>crtN</i> ( <i>crtNa</i> )	ACM07424.1
<i>Halobacillus halophilus</i>	<i>crtNb</i>	ACM07427.1
<i>Halobacillus halophilus</i>	<i>crtNc</i>	ACM07425.1
<i>Methylomonas</i> sp. strain 16a	<i>crtN</i> ( <i>crtNa</i> )	AAX46183.1
<i>Methylomonas</i> sp. strain 16a	<i>crtNb</i>	AAX46185.1
<i>Methylomonas</i> sp. strain 16a	<i>aldH</i> ( <i>crtNc</i> )	AAX46184.1
<i>Planococcus faecalis</i> AJ003 <sup>T</sup>	<i>crtP</i> ( <i>crtP2</i> )	WP_071154552.1
<i>Planococcus maritimus</i> strain iso-3	<i>crtNa</i> , <i>crtNb</i> , <i>orf2</i>	LC620265
<i>Planococcus maritimus</i> strain iso-3	<i>cruO</i> ( <i>crtP</i> )	this study
<i>Rhodobacter capsulatus</i>	<i>crtI</i>	ADE84444.1
<i>Pantoea ananatis</i>	<i>crtI</i>	AER34890
<i>Lactobacillus (Lactiplantibacillus) plantarum</i>	<i>crtN</i> ( <i>crtNa</i> )	WP_011102097.1
<i>Bacillus infantis</i>	<i>crtNb</i>	AGX02538.1
<i>Bacillus (Cytobacillus) firmus</i>	<i>crtNc</i>	AGX02539.1
<i>Bacillus (Cytobacillus) firmus</i>	<i>crtN</i> ( <i>crtNa</i> )	AGX02541.1

### 3.3. *CruO* Acts as an Aldehyde Synthase

To investigate the function of the *cruO* gene from *P. maritimus* iso-3, we constructed plasmids pAC-HIMNFNBp and pAC-HIMNFNBGP (Figure 2) and introduced them into *E. coli* (JM101(DE3)). Recombinant *E. coli* strains carrying plasmids pAC-HIMNFNB and pAC-HIMNFNBG produced 5-hydroxy-5,6-dihydro-apo-4,4'-lycopene (Peak A) and 5-glucosyl-5,6-dihydro-apo-4,4'-lycopene (Peak B), respectively (Figure 4a,c), as shown previously [3]. However, new peaks **1** and **3** were observed in *E. coli* carrying pAC-HIMNFNBp and pAC-HIMNFNBGP, respectively (Figure 4a,c). The produced compounds **1** and **3** were purified and analyzed using ESI-MS (+), <sup>1</sup>H, and <sup>13</sup>C NMR, as described later. Consequently, **1** and **3** were identified as 5-hydroxy-5,6-dihydro-4,4'-diapolycopen-4'-al and 5-glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-al, respectively. Thus, the *cruO* gene was confirmed to encode the carotenoid terminal oxidase, which converts the terminal methyl group of a C<sub>30</sub>-carotenoid into

its aldehydes. In the *crtNb* clade (Figure 3), the *Methylomonas* 16a *crtNb* gene and the *crtP* genes of *S. aureus* and *P. faecalis* are considered to encode enzymes with similar oxidation activity to CruO [15].



**Figure 4.** Functional analysis of *cruO* (*crtP*) and *aldH* of *P. maritimus* strain iso-3. (a) HPLC chromatograms of the extracts of *E. coli* transformants carrying pAC-HIMNFNBp; (b) pAC-HIMNFNBpA; (c) pAC-HIMNFNBGP; and (d) pAC-HIMNFNBGPA. The UV-visible spectra of compounds 1–4 are presented on the right of the chromatograms. Peak A, 5-hydroxy-5,6-dihydro-apo-4,4'-lycopen; peak B, 5-glucosyl-5,6-dihydro-apo-4,4'-lycopen.


### 3.4. AldH Acts as an Aldehyde Dehydrogenase

The *aldH* gene from *P. maritimus* iso-3 is homologous to *aldH* from *P. faecalis*, which encodes an aldehyde dehydrogenase [13]. The catalytic activity of the *P. maritimus* AldH protein was examined by constructing the plasmids pAC-HIMNFNBpA and pAC-HIMNFNBGPA (Figure 2), which were individually introduced into *E. coli* (DE3). These *E. coli* transformants generated new carotenoid peaks 2 and 4, respectively (Figure 4b,d). Their ESI-MS (+),  $^1\text{H}$ , and  $^{13}\text{C}$  NMR spectral analyses indicated compounds 2 and 4 as 5-hydroxy-5,6-dihydro-4,4'-diapolycopen-4'-oic acid and 5-glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-oic acid, respectively. These results show that *aldH* codes for aldehyde dehydrogenase, which converts the terminal aldehyde group of a  $\text{C}_{30}$ -carotenoid into its carboxylic acid.

### 3.5. Structural Determination of the Intermediate Carotenoids (1–4)

The molecular formula of 1 was determined as  $\text{C}_{30}\text{H}_{40}\text{O}_2$  using HR-ESI-MS analysis. Analyses of  $^1\text{H}$  and  $^{13}\text{C}$  NMR,  $^1\text{H}$ - $^1\text{H}$  DQF COSY, and HMQC spectra of 1 in  $\text{CDCl}_3$  showed that the structure of 1 was closely related to that of 4,4'-diapolycopen, while the signals of two singlet methyls (H-4 ( $\delta_{\text{H}}$  1.14) and C-4 ( $\delta_{\text{C}}$  29.2)) and (H-18 ( $\delta_{\text{H}}$  1.14) and C-18 ( $\delta_{\text{C}}$  29.3)), one non-oxygenated  $\text{sp}^3$   $\text{CH}_2$  (H-6 ( $\delta_{\text{H}}$  2.32) and C-6 ( $\delta_{\text{C}}$  47.4)), one oxygenated quaternary  $^{13}\text{C}$  (C-5 ( $\delta_{\text{C}}$  71.0)), and one aldehyde (H-4' ( $\delta_{\text{H}}$  9.45) and C-4' ( $\delta_{\text{C}}$  194.4)) were observed only in 1. The linkages between the observed  $^1\text{H}$  and  $^{13}\text{C}$  signals were analyzed using the HMBC spectrum. In the HMBC spectrum, preservation of the all *trans* olefin structure of 4,4'-diapolycopen from C-7 to C-5 in 1 was proved using the  $^1\text{H}$ - $^{13}\text{C}$  long-range couplings from the methyl signals of H-19 ( $\delta_{\text{H}}$  1.94), H-20 ( $\delta_{\text{H}}$  1.98), H-18' ( $\delta_{\text{H}}$  1.90), H-19' ( $\delta_{\text{H}}$  2.02), and H-20' ( $\delta_{\text{H}}$  1.98) (Figure 5). The  $^1\text{H}$ - $^{13}\text{C}$  long range couplings from H-4 and H-18 to C-5 and C-6 and vicinal  $^1\text{H}$ - $^1\text{H}$  spin coupling between H-6 and H-7 showed that the 2-methyl, 2-oxygenated propyl structure composed of C-4, C-5, C-6, and C-18 were attached at C-7. Furthermore, the aldehyde function at C-4' was demonstrated using the  $^1\text{H}$ - $^{13}\text{C}$  long-range coupling from H-18' to C-4'. Based on these observations, the structure of 1 was determined to be 5-hydroxy-5,6-dihydro-4,4'-diapolycopen-4'-al (Figure 5). According to the CAS database, compound 1 was new.





**Figure 5.** Structural determination of compounds 1–4.

The molecular formula of **2** was determined to be  $\text{C}_{30}\text{H}_{40}\text{O}_3$  by HR-ESI-MS analysis. The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of **2** in  $\text{CDCl}_3$  were almost identical to those of **1** except for the disappearance of the aldehyde signal in **1** and the appearance of a carbonyl signal (C-4' ( $\delta_{\text{C}}$  169.7)). Because long-range coupling from H-18' ( $\delta_{\text{H}}$  1.88) to C-4' was observed in the HMBC spectrum and the molecular formula of **2** was **1** + O, the structure of **2** was determined as 5-hydroxy-5,6-dihydro-4,4'-diapolycopen-4'-oic acid (Figure 5). Compound **2** was a new compound according to the CAS database.

The molecular formula of **3** was determined to be  $\text{C}_{36}\text{H}_{50}\text{O}_7$  using HR-ESIMS analysis. The  $^1\text{H}$  and  $^{13}\text{C}$  NMR,  $^1\text{H}$ - $^1\text{H}$  DQF COSY, and HMQC spectra of **3** in  $\text{CD}_3\text{OD}$  revealed that the structure of **3** was closely related to **1**, while  $^1\text{H}$  and  $^{13}\text{C}$  signals derived from a hexose were observed in **3**. Hexose was identified as  $\beta$ -glucose by  $^1\text{H}$ - $^1\text{H}$  vicinal spin couplings of H-1'-H-6' and the linkage of the glucose at C-5 was shown by the  $^1\text{H}$ - $^{13}\text{C}$  long-range coupling from H-1'' ( $\delta_{\text{H}}$  4.51) to C-5 ( $\delta_{\text{C}}$  70.4) observed in the HMBC spectrum. Based on these findings, the structure of **3** was determined to be 5-glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-al (Figure 5). According to the CAS database, compound **3** was new.

The molecular formula of **4** was determined to be  $\text{C}_{36}\text{H}_{50}\text{O}_7$  using HR-ESI-MS analysis. The  $^1\text{H}$ ,  $^{13}\text{C}$  NMR,  $^1\text{H}$ - $^1\text{H}$  DQF COSY, and HMQC spectra of **4** in  $\text{DMSO}-d_6$  revealed that the structure of **4** was almost identical to **3**, while the aldehyde signal in **3** disappeared in **4** and a carbonyl signal (C-4' ( $\delta_{\text{C}}$  169.5)) was observed only in **4**. Since the long-range coupling from H-18' ( $\delta_{\text{H}}$  1.90) to C-4' was observed in the HMBC spectrum (Figure 5) and the molecular formula of **4** was compound **3** + O, the structure of **4** was determined as 5-glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-oic acid (Figure 5). According to the CAS database, compound **4** was new.

### 3.6. Singlet Oxygen-quenching Activities of Newly Generated Carotenoids

The singlet oxygen-quenching activities of the newly generated C<sub>30</sub>-carotenoids (intermediates) were examined and the results are shown in Table 2. As shown in Table 2, compounds closer to the end of the biosynthetic pathway exhibited more potent singlet oxygen-quenching activity.

**Table 2.** Singlet oxygen-quenching activity of the individual C<sub>30</sub>-carotenoids (intermediates).

Carotenoid	Singlet Oxygen-quenching IC <sub>50</sub> (μM)
15- <i>cis</i> -4,4'-diapophytoene	>100 <sup>a</sup>
4,4'-diaponeurosporene	45 <sup>a</sup>
5-hydroxy-5,6-dihydro-4,4'-diaponeurosporene	56 <sup>a</sup>
5-hydroxy-5,6-dihydro-4,4'-diapolycopene	30 <sup>a</sup>
5-hydroxy-5,6-dihydro-4,4'-diapolycopen-4'-al ( <b>1</b> )	9.3
5-hydroxy-5,6-dihydro-4,4'-diapolycopen-4'-oic acid ( <b>2</b> )	7.8
5-glucosyl-5,6-dihydro-4,4'-diapolycopene	30 <sup>a</sup>
5-glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-al ( <b>3</b> )	8.3
5-glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-oic acid ( <b>4</b> )	12
methyl-5-glucosyl-5,6-dihydro-4,4'-diapolycopenoate	5.1 <sup>a</sup>
astaxanthin	3.7 <sup>a</sup>

<sup>a</sup> cited from the results in Takemura et al. [3] for comparison.

This result suggests that *P. maritimus* iso-3 living near the surface of the sea produces methyl-5-glucosyl-5,6-dihydro-4,4'-diapolycopenoate to protect itself from singlet oxygen damage caused by ultraviolet and triplet oxygen. The dipolar structure of this highly modified C<sub>30</sub>-carotenoid may also stabilize the phospholipid double-layer membrane of *Planococcus*, and consequently may contribute to the tolerance of this bacterium not only to photooxidative damage but also to solvents [3].

A similar highly modified dipolar C<sub>30</sub>-carotenoid, glycosyl-4,4'-diaponeurosporen-4'-ol-4'-oate, is found in the same genus *P. faecalis* [13]. On the other hand, *S. aureus* is known to produce staphyloxanthin as the monopolar C<sub>30</sub>-carotenoid, whose biosynthetic pathway has been elucidated at the gene level [7,8]. We have elucidated the biosynthetic pathway of highly modified dipolar C<sub>30</sub>-carotenoids for the first time, using the *Planococcus* genes.

## 4. Conclusions

In the marine bacterium *Planococcus maritimus* strain iso-3, the *crtP* (here renamed *cruO*) and *aldH* genes were shown to encode carotenoid terminal oxidase (carotenoid-aldehyde synthase) and carotenoid-aldehyde dehydrogenase, respectively. The carotenoid biosynthetic pathway for 5-glucosyl-5,6-dihydro-4,4'-diapolycopen-4'-oic acid (**4**) was identified at the gene level (Figure 1). We further produced four new highly modified C<sub>30</sub>-carotenoids **1–3** in addition to **4**, which were obtained using *E. coli* cells carrying combinations of *P. maritimus* carotenogenic genes. They have been shown to retain potent singlet oxygen-quenching activities. We await further biological evaluations of **1–4** as future studies.

## Supplementary Materials

The following supporting information can be found at: <https://www.sciepublish.com/index/journals/article/sbe/25.html/id/17>.

Data Availability: Accession numbers of the *cruO* (*crtP*) and *aldH* genes are LC722835 and LC722836, respectively. Accession of the sequences of the plasmids, pAC-HIMNFnBP, pAC-HIMNFnbGP, pAC-HIMNFnbPA and pAC-HIMNFnbGPA are LC722947, LC722948, LC722949, and LC722950, respectively.

## Author Contributions

Conceptualization: K.S. and N.M.; Investigation: M. H., C.M., and M.T.; Writing – Original Draft Preparation: K.S., M.T., and N.M.

## Ethics Statement

Not applicable.

## Informed Consent Statement

Not applicable.

## Funding

This research received no external funding.

## Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## References

1. Sandmann G, Misawa N. Carotenoid production in *Escherichia coli*: Case of acyclic carotenoids. In *Carotenoids: Biosynthetic and Biofunctional Approaches*; Misawa N., Ed.; Springer: Singapore, 2021; pp. 201–208.
2. Wieland B, Feil C, Gloria-Maercker E, Thumm G, Lechner M, Bravo JM, et al. Genetic and biochemical analyses of the biosynthesis of the yellow carotenoid 4,4'-diaponeurosporene of *Staphylococcus aureus*. *J. Bacteriol.* **1994**, *176*, 7719–7726.
3. Takemura M, Takagi C, Aikawa M, Araki K, Choi S-K, Itaya M, et al. Heterologous production of novel and rare C<sub>30</sub>-carotenoids using *Planococcus* carotenoid biosynthesis genes. *Microb. Cell Fact.* **2021**, *20*, 194–205.
4. Shindo K, Endo M, Miyake Y, Wakasugi K, Morrith D, Bramley PM, et al. Methyl glucosyl-3,4-dehydro-apo-8'-lycopenoate, a novel antioxidative glyco-C<sub>30</sub>-carotenoic acid produced by a marine bacterium *Planococcus maritimus*. *J. Antibiot.* **2008**, *61*, 729–735.
5. Shindo K, Endo M, Miyake Y, Wakasugi K, Morrith D, Bramley PM, et al. Methyl 5-glucosyl-5,6-dihydro-apo-4,4'-lycopenoate, a novel antioxidative glyco-C<sub>30</sub>-carotenoic acid produced by a marine bacterium *Planococcus maritimus*. *J. Antibiot.* **2014**, *67*, 731–732.
6. Sun Z, Shen S, Wang C, Wang H, Hu Y, Jiao J, et al. A novel carotenoid 1,2-hydratase (CruF) from two species of the non-photosynthetic bacterium *Deinococcus*. *Microbiology* **2009**, *155*, 2775–2783.
7. Kim SH, Lee PC. Functional expression and extension of Staphylococcal staphyloxanthin biosynthesis pathway in *Escherichia coli*. *J. Biol. Chem.* **2012**, *287*, 21575–21583.
8. Pelz A, Wieland KP, Pitzbach K, Hentschel P, Alberl K, Götz F. Structure and biosynthesis of staphyloxanthin from *Staphylococcus aureus*. *J. Biol. Chem.* **2005**, *280*, 32493–32498.
9. Steiger S, Perez-Fons L, Cutting SM, Fraser PD, Sandmann G. Annotation and functional assignment of the genes for the C<sub>30</sub> carotenoid pathways from the genomes of two bacteria: *Bacillus indicus* and *Bacillus firmus*. *Microbiology* **2015**, *161*, 194–202.
10. Nishida Y, Adachi K, Kasai H, Shizuri Y, Shindo K, Sawabe A, et al. Elucidation of a carotenoid biosynthesis gene cluster encoding a novel enzyme, 2,2'-β-hydroxylase, from *Brevundimonas* sp. strain SD212 and combinatorial biosynthesis of new or rare xanthophylls. *Appl. Environ. Microbiol.* **2005**, *71*, 4286–4296.
11. Fraser PD, Pinto MES, Holloway DE, Bramley PM. Application of high-performance liquid chromatography with photodiode array detection to the metabolic profiling of plant isoprenoids. *Plant J.* **2000**, *24*, 551–558.
12. Yokoyama A, Miki W. Composition and presumed biosynthetic pathway of carotenoids in the astaxanthin-producing bacterium *Agrobacterium aurantiacum*. *FEMS Microbiol. Lett.* **1995**, *128*, 139–144.
13. Lee JH, Kim JW, Lee PC. Genome mining reveals two missing CrtP and AldH enzymes in the C<sub>30</sub> carotenoid biosynthesis pathway in *Planococcus faecalis* AJ003<sup>T</sup>. *Molecules* **2020**, *25*, 5892.
14. Harker M, Hirschberg J. Molecular biology of carotenoid biosynthesis in photosynthetic organisms. *Methods Enzymol.* **1998**, *297*, 244–263.
15. Tao L, Schenzle A, Odom JM, Cheng Q. Novel carotenoid oxidase involved in biosynthesis of 4,4'-diapophytoene dialdehyde. *Appl. Environ. Microbiol.* **2005**, *71*, 3294–3301.