


Supplementary Table S1. Mean and Standard Deviations for the ODD Symptoms Used in the Study.

All Participants		
Symptoms	Mean	SD
Temper	2.26	1.53
Touchy	2.48	1.51
Angry	2.09	1.54
Argues	2.31	1.62
Defies	2.21	1.53
Annoys	1.62	1.56
Blames	2.07	1.65


Ranges of scores for all symptoms = 0 to 5.

Supplementary Table S2. Frequencies of Comorbidity Based on Diagnostic Symptom Threshold Number.

No of Disorders	Frequency	Percent	Cumulative Percent
0.00	192	20.2	20.2
1.00	156	16.4	16.4
2.00	136	14.3	14.3
3.00	102	10.7	10.7
4.00	80	8.4	8.4
5.00	84	8.8	8.8
6.00	64	6.7	6.7
7.00	41	4.3	4.3
8.00	36	3.8	3.8
9.00	23	2.4	2.4
10.00	18	1.9	1.9
11.00	7	.7	.7
12.00	7	.7	.7
13.00	1	.1	.1
14.00	1	.1	.1
15.00	3	.3	.3



Supplementary Figure S1. Schematic diagram illustrating the SEM model used to examine the unique associations of ADHDIA with the ODD dimensions.



Supplementary Figure S2. Joint Structural Equation Modelling (SEM) associations of irritability and headstrong with adolescent psychological disorders.